

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
8 February 2007 (08.02.2007)

PCT

(10) International Publication Number
WO 2007/014923 A1

(51) International Patent Classification:

C07D 401/12 (2006.01) A61K 31/538 (2006.01)
C07D 403/12 (2006.01) A61P 31/14 (2006.01)
C07D 413/12 (2006.01)

(21) International Application Number:

PCT/EP2006/064817

(22) International Filing Date: 28 July 2006 (28.07.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
05107071.2 29 July 2005 (29.07.2005) EP(71) Applicants (for all designated States except US): **TIBOTEC PHARMACEUTICALS LTD.** [IE/IE]; Eastgate Village, Eastgate, Little Island, Co Cork (IE). **MEDIVIR AB** [SE/SE]; Lunastigen 7, S-141 44 Huddinge (SE).

(72) Inventors; and



(75) Inventors/Applicants (for US only): **DE KOCK, Herman Augustinus** [BE/BE]; Wolfseind 10, B-2370 Arendonk (BE). **SIMMEN, Kenneth Alan** [GB/BE]; Boterbloemenlaan 35, B-3080 Tervuren (BE). **JÖNSSON, Carl Erik Daniel** [SE/SE]; Medivir AB, Lunastigen 7, S-141 44 Huddinge (SE). **AYESA ALVAREZ, Susana** [ES/SE]; Medivir AB, Lunastigen 7, S-141 44 Huddinge(SE). **CLASSON, Björn Olof** [SE/SE]; Medivir AB, Lunastigen 7, S-141 44 Huddinge. **NILSSON, Karl Magnus** [SE/SE]; Medivir AB, Lunastigen 7, S-141 44 Huddinge (SE). **ROSENQUIST, Åsa Annica Kristina** [SE/SE]; Medivir AB, Lunastigen 7, S-141 44 Huddinge (SE). **SAMUELSSON, Bengt Bertil** [SE/SE]; Medivir AB, Lunastigen 7, S-141 44 Huddinge (SE). **WALLBERG, Hans Kristian** [SE/SE]; Medivir AB, Lunastigen 7, S-141 44 Huddinge (SE).(74) Agent: **DAELEMANS, Frank**; Tibotec-Virco Comm. VA, J & J Patent Law Department, Generaal De Wittelaan L 11B 3, B-2800 Mechelen (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

[Continued on next page]

(54) Title: MACROCYCLIC INHIBITORS OF HEPATITIS C VIRUS



(57) **Abstract:** Inhibitors of HCV replication of formula (I) and the *N*-oxides, salts, or stereoisomers thereof, wherein each dashed line represents an optional double bond; X is N, CH and where X bears a double bond it is C; R¹ is -OR⁶, -NH-SO₂R⁷; R² is hydrogen, and where X is C or CH, R² may also be C₁₋₆alkyl; R³ is hydrogen, C₁₋₆alkyl, C₁₋₆alkoxyC₁₋₆alkyl, or C₃₋₇cycloalkyl; n is 3, 4, 5, or 6; R⁴ and R⁵ taken together with the nitrogen atom to which they are attached form a bicyclic ring system selected from formula (II) wherein said ring system may optionally be substituted with 1-3 substituents; R⁶ is hydrogen; aryl; Het; C₃₋₇cycloalkyl optionally substituted with C₁₋₆alkyl; or C₁₋₆alkyl optionally substituted with C₃₋₇cycloalkyl, aryl or with Het; R⁷ is aryl; Het; C₃₋₇cycloalkyl optionally substituted with C₁₋₆alkyl; or C₁₋₆alkyl optionally substituted with C₃₋₇cycloalkyl, aryl or with Het; aryl is phenyl or naphthyl, each of which may be optionally substituted with 1-3 substituents; Het is a 5 or 6 membered saturated, partially unsaturated or completely unsaturated heterocyclic ring containing 1 to 4 heteroatoms each independently selected from N, O or S, and being optionally substituted with 1-3 substituents pharmaceutical compositions containing compounds (I) and processes for preparing compounds (I). Bioavailable combinations of the inhibitors of HCV of formula (I) with ritonavir are also provided.



GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

MACROCYCLIC INHIBITORS OF HEPATITIS C VIRUS

The present invention is concerned with macrocyclic compounds having inhibitory activity on the replication of the hepatitis C virus (HCV). It further concerns
5 compositions comprising these compounds as active ingredients as well as processes for preparing these compounds and compositions.

Hepatitis C virus is the leading cause of chronic liver disease worldwide and has become a focus of considerable medical research. HCV is a member of the
10 *Flaviviridae* family of viruses in the *hepacivirus* genus, and is closely related to the *flavivirus* genus, which includes a number of viruses implicated in human disease, such as dengue virus and yellow fever virus, and to the animal *pestivirus* family, which includes bovine viral diarrhea virus (BVDV). HCV is a positive-sense, single-stranded
15 RNA virus, with a genome of around 9,600 bases. The genome comprises both 5' and 3' untranslated regions which adopt RNA secondary structures, and a central open reading frame that encodes a single polyprotein of around 3,010-3,030 amino acids. The polyprotein encodes ten gene products which are generated from the precursor polyprotein by an orchestrated series of co- and posttranslational endoproteolytic cleavages mediated by both host and viral proteases. The viral structural proteins
20 include the core nucleocapsid protein, and two envelope glycoproteins E1 and E2. The non-structural (NS) proteins encode some essential viral enzymatic functions (helicase, polymerase, protease), as well as proteins of unknown function. Replication of the viral genome is mediated by an RNA-dependent RNA polymerase, encoded by non-structural protein 5b (NS5B). In addition to the polymerase, the viral helicase and
25 protease functions, both encoded in the bifunctional NS3 protein, have been shown to be essential for replication of HCV RNA. In addition to the NS3 serine protease, HCV also encodes a metalloproteinase in the NS2 region.

Following the initial acute infection, a majority of infected individuals develop chronic
30 hepatitis because HCV replicates preferentially in hepatocytes but is not directly cytopathic. In particular, the lack of a vigorous T-lymphocyte response and the high propensity of the virus to mutate appear to promote a high rate of chronic infection. Chronic hepatitis can progress to liver fibrosis leading to cirrhosis, end-stage liver disease, and HCC (hepatocellular carcinoma), making it the leading cause of liver
35 transplantations.

There are 6 major HCV genotypes and more than 50 subtypes, which are differently distributed geographically. HCV type 1 is the predominant genotype in Europe and the

US. The extensive genetic heterogeneity of HCV has important diagnostic and clinical implications, perhaps explaining difficulties in vaccine development and the lack of response to therapy.

- 5 Transmission of HCV can occur through contact with contaminated blood or blood products, for example following blood transfusion or intravenous drug use. The introduction of diagnostic tests used in blood screening has led to a downward trend in post-transfusion HCV incidence. However, given the slow progression to the end-stage liver disease, the existing infections will continue to present a serious medical and
10 economic burden for decades.

- Current HCV therapies are based on (pegylated) interferon-alpha (IFN- α) in combination with ribavirin. This combination therapy yields a sustained virologic response in more than 40% of patients infected by genotype 1 viruses and about 80% of
15 those infected by genotypes 2 and 3. Beside the limited efficacy on HCV type 1, this combination therapy has significant side effects and is poorly tolerated in many patients. Major side effects include influenza-like symptoms, hematologic abnormalities, and neuropsychiatric symptoms. Hence there is a need for more effective, convenient and better tolerated treatments.

- 20 Recently, two peptidomimetic HCV protease inhibitors have gained attention as clinical candidates, namely BILN-2061 disclosed in WO00/59929 and VX-950 disclosed in WO03/87092. A number of similar HCV protease inhibitors have also been disclosed in the academic and patent literature. It has already become apparent that the sustained
25 administration of BILN-2061 or VX-950 selects HCV mutants which are resistant to the respective drug, so called drug escape mutants. These drug escape mutants have characteristic mutations in the HCV protease genome, notably D168V, D168A and/or A156S. Accordingly, additional drugs with different resistance patterns are required to provide failing patients with treatment options, and combination therapy with multiple
30 drugs is likely to be the norm in the future, even for first line treatment.

- Experience with HIV drugs, and HIV protease inhibitors in particular, has further emphasized that sub-optimal pharmacokinetics and complex dosage regimes quickly result in inadvertent compliance failures. This in turn means that the 24 hour trough
35 concentration (minimum plasma concentration) for the respective drugs in an HIV regime frequently falls below the IC₉₀ or ED₉₀ threshold for large parts of the day. It is considered that a 24 hour trough level of at least the IC₅₀, and more realistically, the IC₉₀ or ED₉₀, is essential to slow down the development of drug escape mutants.

Achieving the necessary pharmacokinetics and drug metabolism to allow such trough levels provides a stringent challenge to drug design. The strong peptidomimetic nature of prior art HCV protease inhibitors, with multiple peptide bonds poses pharmacokinetic hurdles to effective dosage regimes.

5


There is a need for HCV inhibitors which may overcome the disadvantages of current HCV therapy such as side effects, limited efficacy, the emerging of resistance, and compliance failures.

- 10 WO05/037214 relates to macrocyclic carboxylic acids and acylsulfonamides as inhibitors of HCV replication, as well as pharmaceutical compositions, methods of treating a Hepatitis C virus infection and methods of treating liver fibrosis.

The present invention concerns HCV inhibitors which are superior in one or more of
15 the following pharmacological related properties, i.e. potency, decreased cytotoxicity, improved pharmacokinetics, improved resistance profile, acceptable dosage and pill burden.

In addition, the compounds of the present invention have relatively low molecular
20 weight and are easy to synthesize, starting from starting materials that are commercially available or readily available through art-known synthesis procedures.

The present invention concerns inhibitors of HCV replication, which can be represented by formula (I):



25

and the *N*-oxides, salts, and stereoisomers thereof, wherein each dashed line (represented by - - - -) represents an optional double bond;

X is N, CH and where **X** bears a double bond it is C;


R¹ is -OR⁶, -NH-SO₂R⁷;

R² is hydrogen, and where **X** is C or CH, **R**² may also be C₁₋₆alkyl;

R³ is hydrogen, C₁₋₆alkyl, C₁₋₆alkoxyC₁₋₆alkyl, or C₃₋₇cycloalkyl;

5 **n** is 3, 4, 5, or 6;

R⁴ and **R**⁵ taken together with the nitrogen atom to which they are attached form a bicyclic ring system selected from



10 wherein said ring system may optionally be substituted with one, two or three substituents independently selected from halo, hydroxy, oxo, nitro, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₁₋₆alkoxycarbonyl, amino, azido, mercapto, polyhaloC₁₋₆alkyl;

R⁶ is hydrogen; aryl; Het; C₃₋₇cycloalkyl optionally substituted with C₁₋₆alkyl; or C₁₋₆alkyl optionally substituted with C₃₋₇cycloalkyl, aryl or with Het;

15 **R**⁷ is aryl; Het; C₃₋₇cycloalkyl optionally substituted with C₁₋₆alkyl; or C₁₋₆alkyl optionally substituted with C₃₋₇cycloalkyl, aryl or with Het;

aryl as a group or part of a group is phenyl or naphthyl, each of which may be optionally substituted with one, two or three substituents selected from halo, hydroxy, nitro, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkyl, 20 C₁₋₆alkylcarbonyl, amino, mono- or diC₁₋₆alkylamino, azido, mercapto, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkoxy, C₃₋₇cycloalkyl, pyrrolidinyl, piperidinyl, piperazinyl, 4-C₁₋₆alkyl-piperazinyl, 4-C₁₋₆alkylcarbonyl-piperazinyl, and morpholinyl; wherein the morpholinyl and piperidinyl groups may be optionally substituted with one or with two C₁₋₆alkyl radicals;

25 **Het** as a group or part of a group is a 5 or 6 membered saturated, partially unsaturated or completely unsaturated heterocyclic ring containing 1 to 4 heteroatoms each independently selected from nitrogen, oxygen and sulfur, and being optionally substituted with one, two or three substituents each independently selected from the group consisting of halo, hydroxy, nitro, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkoxy, 30 C₁₋₆alkoxyC₁₋₆alkyl, C₁₋₆alkylcarbonyl, amino, mono- or di-C₁₋₆alkylamino, azido, mercapto, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkoxy, C₃₋₇cycloalkyl, pyrrolidinyl, piperidinyl, piperazinyl, 4-C₁₋₆alkyl-piperazinyl, 4-C₁₋₆alkylcarbonyl-piperazinyl,

and morpholinyl; wherein the morpholinyl and piperidinyl groups may be optionally substituted with one or with two C₁₋₆alkyl radicals.

5 The invention further relates to methods for the preparation of the compounds of formula (I), the *N*-oxides, addition salts, quaternary amines, metal complexes, and stereochemically isomeric forms thereof, their intermediates, and the use of the intermediates in the preparation of the compounds of formula (I).

10 The invention relates to the compounds of formula (I) *per se*, the *N*-oxides, addition salts, quaternary amines, metal complexes, and stereochemically isomeric forms thereof, for use as a medicament. The invention further relates to pharmaceutical compositions comprising the aforementioned compounds for administration to a subject suffering from HCV infection. The pharmaceutical compositions may comprise combinations of the aforementioned compounds with other anti-HCV agents.

15 The invention also relates to the use of a compound of formula (I), or a *N*-oxide, addition salt, quaternary amine, metal complex, or stereochemically isomeric forms thereof, for the manufacture of a medicament for inhibiting HCV replication. Or the invention relates to a method of inhibiting HCV replication in a warm-blooded animal
20 said method comprising the administration of an effective amount of a compound of formula (I), or a *N*-oxide, addition salt, quaternary amine, metal complex, or stereochemically isomeric forms thereof.

25 As used in the foregoing and hereinafter, the following definitions apply unless otherwise noted.

The term halo is generic to fluoro, chloro, bromo and iodo.

30 The term "polyhaloC₁₋₆alkyl" as a group or part of a group, e.g. in polyhalo-C₁₋₆alkoxy, is defined as mono- or polyhalo substituted C₁₋₆alkyl, in particular C₁₋₆alkyl substituted with up to one, two, three, four, five, six, or more halo atoms, such as methyl or ethyl with one or more fluoro atoms, for example, difluoromethyl, trifluoromethyl, trifluoroethyl. Preferred is trifluoromethyl. Also included are perfluoroC₁₋₆alkyl groups, which are C₁₋₆alkyl groups wherein all hydrogen atoms are replaced by fluoro atoms, e.g.
35 pentafluoroethyl. In case more than one halogen atom is attached to an alkyl group within the definition of polyhaloC₁₋₆alkyl, the halogen atoms may be the same or different.

As used herein "C₁₋₄alkyl" as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as for example methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-1-propyl;

5 "C₁₋₆alkyl" encompasses C₁₋₄alkyl radicals and the higher homologues thereof having 5 or 6 carbon atoms such as, for example, 1-pentyl, 2-pentyl, 3-pentyl, 1-hexyl, 2-hexyl, 2-methyl-1-butyl, 2-methyl-1-pentyl, 2-ethyl-1-butyl, 3-methyl-2-pentyl, and the like. Of interest amongst C₁₋₆alkyl is C₁₋₄alkyl.

10 The term "C₂₋₆alkenyl" as a group or part of a group defines straight and branched chained hydrocarbon radicals having saturated carbon-carbon bonds and at least one double bond, and having from 2 to 6 carbon atoms, such as, for example, ethenyl (or vinyl), 1-propenyl, 2-propenyl (or allyl), 1-butenyl, 2-butenyl, 3-butenyl, 2-methyl-2-propenyl, 2-pentenyl, 3-pentenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 2-methyl-2-butenyl, 2-methyl-2-pentenyl and the like. Of interest amongst C₂₋₆alkenyl is C₂₋₄alkenyl.

15 The term "C₂₋₆alkynyl" as a group or part of a group defines straight and branched chained hydrocarbon radicals having saturated carbon-carbon bonds and at least one triple bond, and having from 2 to 6 carbon atoms, such as, for example, ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 2-pentyne, 3-pentyne, 20 2-hexynyl, 3-hexynyl and the like. Of interest amongst C₂₋₆alkynyl is C₂₋₄alkynyl.

C₃₋₇cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

25 C₁₋₆alkanediyl defines bivalent straight and branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as, for example, methylene, ethylene, 1,3-propanediyl, 1,4-butanediyl, 1,2-propanediyl, 2,3-butanediyl, 1,5-pentanedyl, 1,6-hexanedyl and the like. Of interest amongst C₁₋₆alkanediyl is C₁₋₄alkanediyl.

30 C₁₋₆alkoxy means C₁₋₆alkyloxy wherein C₁₋₆alkyl is as defined above.

As used herein before, the term (=O) or oxo forms a carbonyl moiety when attached to a carbon atom, a sulfoxide moiety when attached to a sulfur atom and a sulfonyl moiety when two of said terms are attached to a sulfur atom. Whenever a ring or ring system
35 is substituted with an oxo group, the carbon atom to which the oxo is linked is a saturated carbon.

The radical Het is a heterocycle as specified in this specification and claims. Examples of Het comprise, for example, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyrrolyl, imidazolyl, oxazolyl, isoxazolyl, thiazinyl, isothiazinyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl (including 1,2,3-triazolyl, 1,2,4-triazolyl), tetrazolyl, furanyl, thienyl, pyridyl, pyrimidyl, pyridazinyl, pyrazolyl, triazinyl, and the like. Of interest amongst the Het radicals are those which are non-saturated, in particular those having an aromatic character. Of further interest are those Het radicals having one or two nitrogens.

- Each of the Het radicals mentioned in this and the following paragraphs may be optionally substituted with the number and kind of substituents mentioned in the definitions of the compounds of formula (I) or any of the subgroups of compounds of formula (I). Some of the Het radicals mentioned in this and the following paragraphs may be substituted with one, two or three hydroxy substituents. Such hydroxy substituted rings may occur as their tautomeric forms bearing keto groups. For example a 3-hydroxypyridazine moiety can occur in its tautomeric form 2*H*-pyridazin-3-one. Where Het is piperazinyl, it preferably is substituted in its 4-position by a substituent linked to the 4-nitrogen with a carbon atom, e.g. 4-C₁₋₆alkyl, 4-polyhalo-C₁₋₆alkyl, C₁₋₆alkoxyC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl.

- Interesting Het radicals comprise, for example pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl (including 1,2,3-triazolyl, 1,2,4-triazolyl), tetrazolyl, furanyl, thienyl, pyridyl, pyrimidyl, pyridazinyl, pyrazolyl, triazinyl, or any of such heterocycles condensed with a benzene ring, such as indolyl, indazolyl (in particular 1*H*-indazolyl), indolinyl, quinolinyl, tetrahydroquinolinyl (in particular 1,2,3,4-tetrahydroquinolinyl), isoquinolinyl, tetrahydroisoquinolinyl (in particular 1,2,3,4-tetrahydroisoquinolinyl), quinazolinyl, phthalazinyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzoxadiazolyl, benzothiadiazolyl, benzofuranyl, benzothienyl.

- The Het radicals pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, 4-substituted piperazinyl preferably are linked via their nitrogen atom (i.e. 1-pyrrolidinyl, 1-piperidinyl, 4-thiomorpholinyl, 4-morpholinyl, 1-piperazinyl, 4-substituted 1-piperazinyl).

It should be noted that the radical positions on any molecular moiety used in the definitions may be anywhere on such moiety as long as it is chemically stable.

Radicals used in the definitions of the variables include all possible isomers unless otherwise indicated. For instance pyridyl includes 2-pyridyl, 3-pyridyl and 4-pyridyl; pentyl includes 1-pentyl, 2-pentyl and 3-pentyl.

5

When any variable occurs more than one time in any constituent, each definition is independent.

Whenever used hereinafter, the term “compounds of formula (I)”, or “the present compounds” or similar terms, it is meant to include the compounds of formula (I), each and any of the subgroups thereof, their prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes, and stereochemically isomeric forms. One embodiment comprises the compounds of formula (I) or any subgroup of compounds of formula (I) specified herein, as well as the *N*-oxides, salts, as the possible stereoisomeric forms thereof. Another embodiment comprises the compounds of formula (I) or any subgroup of compounds of formula (I) specified herein, as well as the salts as the possible stereoisomeric forms thereof.

The compounds of formula (I) have several centers of chirality and exist as stereochemically isomeric forms. The term “stereochemically isomeric forms” as used herein defines all the possible compounds made up of the same atoms bonded by the same sequence of bonds but having different three-dimensional structures which are not interchangeable, which the compounds of formula (I) may possess.

With reference to the instances where (*R*) or (*S*) is used to designate the absolute configuration of a chiral atom within a substituent, the designation is done taking into consideration the whole compound and not the substituent in isolation.

Unless otherwise mentioned or indicated, the chemical designation of a compound encompasses the mixture of all possible stereochemically isomeric forms, which said compound may possess. Said mixture may contain all diastereomers and/or enantiomers of the basic molecular structure of said compound. All stereochemically isomeric forms of the compounds of the present invention both in pure form or mixed with each other are intended to be embraced within the scope of the present invention.

35

Pure stereoisomeric forms of the compounds and intermediates as mentioned herein are defined as isomers substantially free of other enantiomeric or diastereomeric forms of the same basic molecular structure of said compounds or intermediates. In particular,

the term "stereoisomerically pure" concerns compounds or intermediates having a stereoisomeric excess of at least 80% (i.e. minimum 90% of one isomer and maximum 10% of the other possible isomers) up to a stereoisomeric excess of 100% (i.e. 100% of one isomer and none of the other), more in particular, compounds or intermediates
5 having a stereoisomeric excess of 90% up to 100%, even more in particular having a stereoisomeric excess of 94% up to 100% and most in particular having a stereoisomeric excess of 97% up to 100%. The terms "enantiomerically pure" and "diastereomerically pure" should be understood in a similar way, but then having regard to the enantiomeric excess, and the diastereomeric excess, respectively, of the
10 mixture in question.

Pure stereoisomeric forms of the compounds and intermediates of this invention may be obtained by the application of art-known procedures. For instance, enantiomers may be separated from each other by the selective crystallization of their diastereomeric
15 salts with optically active acids or bases. Examples thereof are tartaric acid, dibenzoyltartaric acid, ditoluoyltartaric acid and camphosulfonic acid. Alternatively, enantiomers may be separated by chromatographic techniques using chiral stationary phases. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting
20 materials, provided that the reaction occurs stereospecifically. Preferably, if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

25 The diastereomeric racemates of the compounds of formula (I) can be obtained separately by conventional methods. Appropriate physical separation methods that may advantageously be employed are, for example, selective crystallization and chromatography, e.g. column chromatography.

30 For some of the compounds of formula (I), their prodrugs, *N*-oxides, salts, solvates, quaternary amines, or metal complexes, and the intermediates used in the preparation thereof, the absolute stereochemical configuration was not experimentally determined. A person skilled in the art is able to determine the absolute configuration of such compounds using art-known methods such as, for example, X-ray diffraction.

35

The present invention is also intended to include all isotopes of atoms occurring on the present compounds. Isotopes include those atoms having the same atomic number but

different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.

The present invention is also intended to include prodrugs of the compounds of formula (I). The term “prodrug” as used throughout this text means the pharmacologically acceptable derivatives such as esters, amides and phosphates, such that the resulting *in vivo* biotransformation product of the derivative is the active drug as defined in the compounds of formula (I). The reference by Goodman and Gilman (The Pharmacological Basis of Therapeutics, 8th ed, McGraw-Hill, Int. Ed. 1992, “Biotransformation of Drugs”, p 13–15) describing prodrugs generally is hereby incorporated. Prodrugs preferably have excellent aqueous solubility, increased bioavailability and are readily metabolized into the active inhibitors *in vivo*. Prodrugs of a compound of the present invention may be prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either by routine manipulation or *in vivo*, to the parent compound.

Preferred are pharmaceutically acceptable ester prodrugs that are hydrolysable *in vivo* and are derived from those compounds of formula (I) having a hydroxy or a carboxyl group. An *in vivo* hydrolysable ester is an ester, which is hydrolyzed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically acceptable esters for carboxy include C₁₋₆alkoxymethyl esters for example methoxymethyl, C₁₋₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl which may be formed at any carboxy group in the compounds of this invention.

An *in vivo* hydrolysable ester of a compound of the formula (I) containing a hydroxy group includes inorganic esters such as phosphate esters and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxy-methoxy. A selection of *in vivo* hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and N-(dialkylaminoethyl)-N-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl. Examples of substituents on

benzoyl include morpholino and piperazino linked from a ring nitrogen atom via a methylene group to the 3- or 4-position of the benzoyl ring.

For therapeutic use, salts of the compounds of formula (I) are those wherein the counter-ion is pharmaceutically acceptable. However, salts of acids and bases which are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

The pharmaceutically acceptable acid and base addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid and base addition salt forms which the compounds of formula (I) are able to form. The pharmaceutically acceptable acid addition salts can conveniently be obtained by treating the base form with such appropriate acid. Appropriate acids comprise, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid, sulfuric, nitric, phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic (i.e. ethanedioic), malonic, succinic (i.e. butanedioic acid), maleic, fumaric, malic (i.e. hydroxybutanedioic acid), tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

Conversely said salt forms can be converted by treatment with an appropriate base into the free base form.

The compounds of formula (I) containing an acidic proton may also be converted into their non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.


The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of formula (I) are able to form by reaction between a basic nitrogen of a compound of formula (I) and an appropriate quaternizing agent, such as, for example, an optionally substituted alkylhalide, arylhalide or arylalkylhalide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl *p*-toluenesulfonates. A quaternary amine has a positively charged nitrogen. Pharmaceutically acceptable counterions include chloro, bromo, iodo, trifluoroacetate and acetate. The counterion of choice can be introduced using ion exchange resins.

The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide.

It will be appreciated that the compounds of formula (I) may have metal binding, chelating, complex forming properties and therefore may exist as metal complexes or metal chelates. Such metalated derivatives of the compounds of formula (I) are intended to be included within the scope of the present invention.

Some of the compounds of formula (I) may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

As mentioned above, the compounds of formula (I) have several asymmetric centers. In order to more efficiently refer to each of these asymmetric centers, the numbering system as indicated in the following structural formula will be used.




Asymmetric centers are present at positions 1, 4 and 6 of the macrocycle as well as at the carbon atom 3' in the 5-membered ring, carbon atom 2' when the R² substituent is C₁₋₆alkyl, and at carbon atom 1' when X is CH. Each of these asymmetric centers can occur in their R or S configuration.

5


The stereochemistry at position 1 preferably corresponds to that of an L-amino acid configuration, i.e. that of L-proline.

10

When X is CH, the 2 carbonyl groups substituted at positions 1' and 5' of the cyclopentane ring preferably are in a trans configuration. The carbonyl substituent at position 5' preferably is in that configuration that corresponds to an L-proline configuration. The carbonyl groups substituted at positions 1' and 5' preferably are as depicted below in the structure of the following formula



15 The compounds of formula (I) include a cyclopropyl group as represented in the structural fragment below:




wherein C₇ represents the carbon at position 7 and carbons at position 4 and 6 are asymmetric carbon atoms of the cyclopropane ring.


20

Notwithstanding other possible asymmetric centers at other segments of the compounds of formula (I), the presence of these two asymmetric centers means that the compounds can exist as mixtures of diastereomers, such as the diastereomers of compounds of formula (I) wherein the carbon at position 7 is configured either syn to the carbonyl or syn to the amide as shown below.


25



- One embodiment concerns compounds of formula (I) wherein the carbon at position 7 is configured syn to the carbonyl. Another embodiment concerns compounds of formula (I) wherein the configuration at the carbon at position 4 is R. A specific subgroup of compounds of formula (I) are those wherein the carbon at position 7 is configured syn to the carbonyl and wherein the configuration at the carbon at position 4 is R.
- 10 The compounds of formula (I) may include a proline residue (when X is N) or a cyclopentyl or cyclopentenyl residue (when X is CH or C).. Preferred are the compounds of formula (I) wherein the substituent at the 1 (or 5') position and the carbamate substituent at position 3' are in a trans configuration. Of particular interest are the compounds of formula (I) wherein position 1 has the configuration
- 15 corresponding to L-proline and the carbamate substituent at position 3' is in a trans configuration in respect of position 1. Preferably the compounds of formula (I) have the stereochemistry as indicated in the structures of formulae (I-a) and (I-b) below:



(I-a)



(I-b)

One embodiment of the present invention concerns compounds of formula (I) or of
 5 formula (I-a) or of any subgroup of compounds of formula (I), wherein one or more of the following conditions apply:


- (a) R^2 is hydrogen;
- (b) X is nitrogen;
- (c) a double bond is present between carbon atoms 7 and 8.

10


One embodiment of the present invention concerns compounds of formula (I) or of
 formulae (I-a), (I-b), or of any subgroup of compounds of formula (I), wherein one or
 more of the following conditions apply:

- (a) R^2 is hydrogen;
- 15 (b) X is CH;
- (c) a double bond is present between carbon atoms 7 and 8.

Particular subgroups of compounds of formula (I) are those represented by the
 following structural formulae:



(I-c)




(I-d)

Amongst the compounds of formula (I-c) and (I-d), those having the stereochemical configuration of the compounds of formulae (I-a), and (I-b), respectively, are of particular interest.


The double bond between carbon atoms 7 and 8 in the compounds of formula (I), or in any subgroup of compounds of formula (I), may be in a *cis* or in a *trans* configuration. Preferably the double bond between carbon atoms 7 and 8 is in a *cis* configuration, as depicted in formulae (I-c) and (I-d).

A double bond between carbon atoms 1' and 2' may be present in the compounds of formula (I), or in any subgroup of compounds of formula (I), as depicted in formula (I-e) below.




(I-e)


Yet another particular subgroup of compounds of formula (I) are those represented by the following structural formulae:



(I-f)



(I-g)



(I-h)

Amongst the compounds of formulae (I-f), (I-g) or (I-h), those having the stereochemical configuration of the compounds of formulae (I-a) and (I-b) are of particular interest.

In (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g) and (I-h), where applicable, X, n, R¹, R², R³, R⁴ and R⁵ are as specified in the definitions of the compounds of formula (I) or in any of the subgroups of compounds of formula (I) specified herein.

It is to be understood that the above defined subgroups of compounds of formulae (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g) or (I-h), as well as any other subgroup defined herein, are meant to also comprise any prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms of such compounds.

5

When *n* is 2, the moiety –CH₂– bracketed by “*n*” corresponds to ethanediyl in the compounds of formula (I) or in any subgroup of compounds of formula (I). When *n* is 3, the moiety –CH₂– bracketed by “*n*” corresponds to propanediyl in the compounds of formula (I) or in any subgroup of compounds of formula (I). When *n* is 4, the moiety
10 –CH₂– bracketed by “*n*” corresponds to butanediyl in the compounds of formula (I) or in any subgroup of compounds of formula (I). When *n* is 5, the moiety –CH₂– bracketed by “*n*” corresponds to pentanediyl in the compounds of formula (I) or in any subgroup of compounds of formula (I). When *n* is 6, the moiety –CH₂– bracketed by “*n*” corresponds to hexanediyl in the compounds of formula (I) or in any subgroup of
15 compounds of formula (I). Particular subgroups of the compounds of formula (I) are those compounds wherein *n* is 4 or 5.

Embodiments of the invention are compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein

- 20 (a) R¹ is –OR⁶, in particular wherein R⁶ is C₁₋₆alkyl, such as methyl, ethyl, or *tert*-butyl and most preferably where R⁶ is hydrogen; or
(b) R¹ is –NHS(=O)₂R⁷, in particular wherein R⁷ is C₁₋₆alkyl, C₃₋₇cycloalkyl, or aryl, e.g. wherein R⁷ is methyl, cyclopropyl, or phenyl; or
(c) R¹ is –NHS(=O)₂R⁷, in particular wherein R⁷ is C₃₋₇cycloalkyl substituted with
25 C₁₋₆alkyl, preferably wherein R⁷ is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl, any of which is substituted with C₁₋₄alkyl, i.e. with methyl, ethyl, propyl, isopropyl, butyl, *tert*-butyl, or isobutyl.

Further embodiments of the invention are compounds of formula (I) or any of the
30 subgroups of compounds of formula (I) wherein R¹ is –NHS(=O)₂R⁷, in particular wherein R⁷ is cyclopropyl substituted with C₁₋₄alkyl, i.e. with methyl, ethyl, propyl, or isopropyl.

Further embodiments of the invention are compounds of formula (I) or any of the
35 subgroups of compounds of formula (I) wherein R¹ is –NHS(=O)₂R⁷, in particular wherein R⁷ is 1-methylcyclopropyl.

Further embodiments of the invention are compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein

- (a) R^2 is hydrogen;
- (b) R^2 is C_{1-6} alkyl, preferably methyl.

5

Embodiments of the invention are compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein

- (a) X is N, C (X being linked via a double bond) or CH (X being linked via a single bond) and R^2 is hydrogen;
- 10 (b) X is C (X being linked via a double bond) and R^2 is C_{1-6} alkyl, preferably methyl.

Further embodiments of the invention are compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein


- (a) R^3 is hydrogen;
- 15 (b) R^3 is C_{1-6} alkyl;
- (d) R^3 is C_{1-6} alkoxy C_{1-6} alkyl or C_{3-7} cycloalkyl.

Preferred embodiments of the invention are compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein R^3 is hydrogen, or C_{1-6} alkyl, more preferably hydrogen or methyl.

20

Embodiments of the invention are compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein R^4 and R^5 taken together with the nitrogen atom to which they are attached form a bicyclic ring system selected from


25



wherein said ring system may optionally be substituted with one or two substituents independently selected from halo, hydroxy, oxo, cyano, carboxyl, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkoxy C_{1-6} alkyl, C_{1-6} alkoxycarbonyl, amino, and polyhalo C_{1-6} alkyl.


- 30 Other subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein R^4 and R^5 taken

together with the nitrogen atom to which they are attached form a bicyclic ring system selected from




- wherein said ring system may optionally be substituted with one or two substituents
 5 independently selected from fluoro, chloro, hydroxy, oxo, cyano, carboxyl, methyl, ethyl, isopropyl, t-butyl, methoxy, ethoxy, isopropoxy, *tert*-butoxy, methoxyethyl, ethoxymethyl, methoxycarbonyl, ethoxycarbonyl, amino, and trifluoromethyl.


- Embodiments of the invention are compounds of formula (I) or any of the subgroups of
 10 compounds of formula (I) wherein R^4 and R^5 taken together with the nitrogen atom to which they are attached form a bicyclic ring system selected from



- wherein the phenyl of said bicyclic ring system is optionally substituted with one or
 two substituents independently selected from halo, hydroxy, cyano, carboxyl, C_{1-6} alkyl,
 15 C_{1-6} alkoxy, C_{1-6} alkoxy-carbonyl, amino, and polyhalo C_{1-6} alkyl.


Embodiments of the invention are compounds of formula (I) or any of the subgroups of
 compounds of formula (I) wherein R^4 and R^5 taken together with the nitrogen atom to
 which they are attached form a bicyclic ring system selected from






wherein said bicyclic ring system is optionally substituted on the phenyl part,
preferably at the positions above indicated with dotted lines with one or two
substituents independently selected from halo, hydroxy, cyano, carboxyl, C₁₋₆alkyl,
5 C₁₋₆alkoxy, C₁₋₆alkoxy-carbonyl, amino, and polyhalo-C₁₋₆alkyl.

Embodiments of the invention are compounds of formula (I) or any of the subgroups of
compounds of formula (I) wherein R⁴ and R⁵ taken together with the nitrogen atom to
which they are attached form a bicyclic ring system selected from



wherein the pyrrolidine, piperidine, or morpholine rings of said bicyclic ring system are
optionally substituted with one or two substituents independently selected from
C₁₋₆alkyl, C₁₋₆alkoxy, and C₁₋₆alkoxyC₁₋₆alkyl.

15 Embodiments of the invention are compounds of formula (I) or any of the subgroups of
compounds of formula (I) wherein R⁴ and R⁵ taken together with the nitrogen atom to
which they are attached form a bicyclic ring system selected from




20 wherein the phenyl of said bicyclic ring system is optionally substituted with one
substituent independently selected from halo, hydroxy, cyano, carboxyl, C₁₋₆alkyl,

C₁₋₆alkoxy, C₁₋₆alkoxy-carbonyl, amino, and polyhaloC₁₋₆alkyl; and wherein the pyrrolidine, piperidine, or morpholine rings of said bicyclic ring system are optionally substituted with one substituent independently selected from C₁₋₆alkyl, C₁₋₆alkoxy, and C₁₋₆alkoxyC₁₋₆alkyl.

5


Embodiments of the invention are compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein R⁴ and R⁵ taken together with the nitrogen atom to which they are attached form a bicyclic ring system selected from



10

wherein said bicyclic ring system is optionally substituted on the positions above indicated with one or two substituents independently selected from C₁₋₆alkyl, C₁₋₆alkoxy, and C₁₋₆alkoxyC₁₋₆alkyl.

15 Preferred embodiments of the invention are compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein R⁴ and R⁵ taken together with the nitrogen atom to which they are attached form a bicyclic ring system selected from





20 The compounds of formula (I) consist of three building blocks P1, P2, P3. Building block P1 further contains a P1' tail. The carbonyl group marked with an asterisk in compound (I-c) below may be part of either building block P2 or of building block P3.

For reasons of chemistry, building block P2 of the compounds of formula (I) wherein X is C incorporates the carbonyl group attached to the position 1'.

The linking of building blocks P1 with P2, P2 with P3, and P1 with P1' (when R¹ is -NH-SO₂R⁷) involves forming an amide bond. The linking of blocks P1 and P3 involves double bond formation. The linking of building blocks P1, P2 and P3 to prepare compounds (I-i) or (I-j) can be done in any given sequence. One of the steps involves a cyclization whereby the macrocycle is formed.

10 Represented herebelow are compounds (I-i) which are compounds of formula (I) wherein carbon atoms C7 and C8 are linked by a double bond, and compounds (I-j) which are compounds of formula (I) wherein carbon atoms C7 and C8 are linked by a single bond. The compounds of formula (I-j) can be prepared from the corresponding compounds of formula (I-I) by reducing the double bond in the macrocycle.


15



20 The synthesis procedures described hereinafter are meant to be applicable for as well the racemates, stereochemically pure intermediates or end products, as any stereoisomeric mixtures. The racemates or stereochemical mixtures may be separated into stereoisomeric forms at any stage of the synthesis procedures. In one embodiment, the intermediates and end products have the stereochemistry specified above in the compounds of formula (I-a) and (I-b).

25


In the synthesis procedures described hereinafter, R^8 represents a radical



wherein the dotted line represents the bond by which the radical is linked to the remainder of the molecule.

5

In a preferred embodiment, compounds (I) wherein the bond between C_7 and C_8 is a double bond, which are compounds of formula (I-i), as defined above, may be prepared as outlined in the following reaction scheme:



- 10 Formation of the macrocycle can be carried out via an olefin metathesis reaction in the presence of a suitable metal catalyst such as e.g. the Ru-based catalyst reported by Miller, S.J., Blackwell, H.E., Grubbs, R.H. J. Am. Chem. Soc. 118, (1996), 9606-9614; Kingsbury, J. S., Harrity, J. P. A., Bonitatebus, P. J., Hoveyda, A. H., J. Am. Chem. Soc. 121, (1999), 791-799; and Huang et al., J. Am. Chem. Soc. 121, (1999), 2674-
15 2678; for example a Hoveyda-Grubbs catalyst.

- Air-stable ruthenium catalysts such as bis(tricyclohexylphosphine)-3-phenyl-1H-inden-1-ylidene ruthenium chloride (Neolyst M1[®]) or bis(tricyclohexylphosphine)-[(phenylthio)methylene]ruthenium (IV) dichloride can be used. Other catalysts that can
20 be used are Grubbs first and second generation catalysts, i.e. Benzyldiene-bis(tricyclohexylphosphine)dichlororuthenium and (1,3-bis-(2,4,6-trimethylphenyl)-2-imidazolidinylidene)dichloro(phenylmethylene)-(tricyclohexylphosphine)ruthenium, respectively. Of particular interest are the Hoveyda-Grubbs first and second generation catalysts, which are dichloro(*o*-isopropoxyphenylmethylene)(tricyclohexylphosphine)-
25 ruthenium(II) and 1,3-bis-(2,4,6-trimethylphenyl)-2-imidazolidinylidene)dichloro-


(*o*-isopropoxyphenylmethylene)ruthenium respectively. Also other catalysts containing other transition metals such as Mo can be used for this reaction.


The metathesis reactions may be conducted in a suitable solvent such as for example ethers, e.g. THF, dioxane; halogenated hydrocarbons, e.g. dichloromethane, CHCl_3 , 1,2-dichloroethane and the like, hydrocarbons, e.g. toluene. In a preferred embodiment, the metathesis reaction is conducted in toluene. These reactions are conducted at increased temperatures under nitrogen atmosphere.

10 Compounds of formula (I) wherein the link between C7 and C8 in the macrocycle is a single bond, i.e. compounds of formula (I-j), can be prepared from the compounds of formula (I-i) by a reduction of the C7-C8 double bond in the compounds of formula (I-i). This reduction may be conducted by catalytic hydrogenation with hydrogen in the presence of a noble metal catalyst such as, for example, Pt, Pd, Rh, Ru or Raney nickel.

15 Of interest is Rh on alumina. The hydrogenation reaction preferably is conducted in a solvent such as, e.g. an alcohol such as methanol, ethanol, or an ether such as THF, or mixtures thereof. Water can also be added to these solvents or solvent mixtures.


The R¹ group can be connected to the P1 building block at any stage of the synthesis, i.e. before or after the cyclization, or before or after the cyclization and reduction as described herein above. The compounds of formula (I) wherein R¹ represents -NHSO₂R⁷, said compounds being represented by formula (I-k-1), can be prepared by linking the R¹ group to P1 by forming an amide bond between both moieties. Similarly, the compounds of formula (I) wherein R¹ represents -OR⁶, i.e. compounds (I-k-2), can be prepared by linking the R¹ group to P1 by forming an ester bond. In one embodiment, the -OR⁶ groups are introduced in the last step of the synthesis of the compounds (I) as outlined in the following reaction schemes wherein G represents a group:





Intermediate (2a) can be coupled with the amine (2b) by an amide forming reaction such as any of the procedures for the formation of an amide bond described hereinafter. In particular, (2a) may be treated with a coupling agent, for example *N,N'*-carbonyl-diimidazole (CDI), EEDQ, IIDQ, EDCI or benzotriazol-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate (commercially available as PyBOP®), in a solvent such as an ether, e.g. THF, or a halogenated hydrocarbon, e.g. dichloromethane, chloroform, dichloroethane, and reacted with the desired sulfonamide (2b), preferably after reacting (2a) with the coupling agent. The reactions of (2a) with (2b) preferably are conducted in the presence of a base, for example a trialkylamine such as triethylamine or diisopropylethylamine, or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). Intermediate (2a) can also be converted into an activated form, e.g. an activated form of general formula G-CO-Z, wherein Z represents halo, or the rest of an active ester, e.g. Z is an aryloxy group such as phenoxy, p-nitrophenoxy, pentafluorophenoxy, trichlorophenoxy, pentachlorophenoxy and the like; or Z can be the rest of a mixed anhydride. In one embodiment, G-CO-Z is an acid chloride (G-CO-Cl) or a mixed acid anhydride (G-CO-O-CO-R or G-CO-O-CO-OR, R in the latter being e.g. C₁₋₄alkyl, such as methyl, ethyl, propyl, i-propyl, butyl, t-butyl, i-butyl, or benzyl). The activated form G-CO-Z is reacted with the sulfonamide (2b).


The activation of the carboxylic acid in (2a) as described in the above reactions may lead to an internal cyclization reaction to an azalactone intermediate of formula



wherein X, R², R³, R⁴, R⁵, n are as specified above and wherein the stereogenic centers may have the stereochemical configuration as specified above, for example as in (I-a) or (I-b). The intermediates (2a-1) can be isolated from the reaction mixture, using
 5 conventional methodology, and the isolated intermediate (2a-1) is then reacted with (2b), or the reaction mixture containing (2a-1) can be reacted further with (2b) without isolation of (2a-1). In one embodiment, where the reaction with the coupling agent is conducted in a water-immiscible solvent, the reaction mixture containing (2a-1) may be washed with water or with slightly basic water in order to remove all water-soluble side
 10 products. The thus obtained washed solution may then be reacted with (2b) without additional purification steps. The isolation of intermediates (2a-1) on the other hand may provide certain advantages in that the isolated product, after optional further purification, may be reacted with (2b), giving rise to less side products and an easier work-up of the reaction.


15 Intermediate (2a) can be coupled with the alcohol (2c) by an ester forming reaction. For example, (2a) and (2c) are reacted together with removal of water either physically, e.g. by azeotropic water removal, or chemically by using a dehydrating agent. Intermediate (2a) can also be converted into an activated form G-CO-Z, such as the
 20 activated forms mentioned above, and subsequently reacted with the alcohol (2c). The ester forming reactions preferably are conducted in the presence of a base such as an alkali metal carbonate or hydrogen carbonate, e.g. sodium or potassium hydrogen carbonate, or a tertiary amine such as the amines mentioned herein in relation to the amide forming reactions, in particular a trialkylamine, e.g. triethylamine. Solvents that
 25 can be used in the ester forming reactions comprise ethers such as THF; halogenated hydrocarbons such as dichloromethane, CH₂Cl₂; hydrocarbons such as toluene; polar aprotic solvents such as DMF, DMSO, DMA; and the like solvents.

The compounds of formula (I) wherein R^3 is hydrogen, said compounds being represented by (I-I), can also be prepared by removal of a protecting group PG, from a corresponding nitrogen-protected intermediate (3a), as in the following reaction scheme. The protecting group PG in particular is any of the nitrogen protecting groups mentioned hereinafter and can be removed using procedures also mentioned hereinafter:



The starting materials (3a) in the above reaction can be prepared following the procedures for the preparation of compounds of formula (I), but using intermediates wherein the group R^3 is PG.


The compounds of formula (I) can also be prepared by reacting an intermediate (4a) with an amine (4b) in the presence of a carbamate forming reagent as outlined in the following reaction scheme wherein the various radicals have the meanings specified above:



The reaction of intermediates (4a) with the carbamate forming reagent is conducted in the same solvents and bases as those used for the amide bond formation as described hereinafter.

Carbamate forming reactions may be conducted using a variety of methods, in particular by reaction of amines with alkyl chloroformates; by reaction of alcohols with carbamoyl chlorides or isocyanates; via reactions involving metal complexes or acyl transfer agents. See for example, Greene, T. W. and Wuts, P. G. M., "Protective Groups in Organic Synthesis"; 1999; Wiley and Sons, p. 309-348. Carbon monoxide and certain metal catalysts can be used to synthesize carbamates from several starting compounds, including amines. Metals such as palladium, iridium, uranium, and platinum may be used as catalysts. Methods using carbon dioxide for synthesis of carbamates that have been also been reported, can also be used (see for example, Yoshida, Y., et al., *Bull. Chem. Soc.* Japan 1989, 62, 1534; and Aresta, M., et al., *Tetrahedron*, 1991, 47, 9489).

One approach for the preparation of carbamates involves the use of intermediates



15


wherein Q is leaving group such as halo, in particular chloro and bromo, or a group used in active esters for amide bond formation, such as those mentioned above, for example phenoxy or substituted phenoxy such as p.chloro and p.nitrophenoxy, trichlorophenoxy, pentachlorophenoxy, N-hydroxy-succinimidyl, and the like.

20


Intermediates (4b) can be derived from alcohols (4a) and phosgene, thus forming a chloroformate, or by transferring the chloro in the latter to intermediates (5a) which are intermediates of formula (5) wherein Q is Q¹. In this and the following reaction procedures, Q¹ represents any of the active ester moieties such as those mentioned above. Intermediates (4b) are reacted with (4a), obtaining compounds (I).

25

Intermediates (4b-1), which are intermediates (4b) wherein Q is Q¹, can also be prepared by reacting the alcohol (4a) with carbonates Q¹-CO-Q¹ such as e.g. bisphenol, bis-(substituted phenol) or bis N-hydroxy-succinimidyl carbonates:



The reagents (5a) may also be prepared from chloroformates $Cl-CO-Q^1$ as follows:



5 The above reactions to prepare reagents (4b-1) may be conducted in the presence of the bases and solvents mentioned hereinafter for the synthesis of amide bonds, in particular triethylamine and dichloromethane.

10 Alternatively, in order to prepare the compounds of formula (I), first an amide bond between building blocks P2 and P1 is formed, followed by coupling of the P3 building block to the P1 moiety in P1-P2, and a subsequent carbamate or ester bond formation between P3 and the P2 moiety in P2-P1-P3 with concomitant ring closure.

15 Yet another alternative synthetic methodology is the formation of an amide bond between building blocks P2 and P3, followed by the coupling of building block P1 to the P3 moiety in P3-P2, and a last amide bond formation between P1 and P2 in P1-P3-P2 with concomitant ring closure.

20 Building blocks P1 and P3 can be linked to a P1-P3 sequence. If desired, the double bond linking P1 and P3 may be reduced. The thus formed P1-P3 sequence, either reduced or not, can be coupled to building block P2 and the thus forming sequence P1-P3-P2 subsequently cyclized, by forming an amide bond.

25 Building blocks P1 and P3 in any of the previous approaches can be linked via double bond formation, e.g. by the olefin metathesis reaction described hereinafter, or a Wittig type reaction. If desired, the thus formed double bond can be reduced, similarly as described above for the conversion of (I-i) to (I-j). The double bond can also be reduced at a later stage, i.e. after addition of a third building block, or after formation of

the macrocycle. Building blocks P2 and P1 are linked by amide bond formation and P3 and P2 are linked by carbamate or ester formation.

5 The tail P1' can be bonded to the P1 building block at any stage of the synthesis of the compounds of formula (I), for example before or after coupling the building blocks P2 and P1; before or after coupling the P3 building block to P1; or before or after ring closure.

10 The individual building blocks can first be prepared and subsequently coupled together or alternatively, precursors of the building blocks can be coupled together and modified at a later stage to the desired molecular composition.

15 The formation of amide bonds can be carried out using standard procedures such as those used for coupling amino acids in peptide synthesis. The latter involves the dehydrative coupling of a carboxyl group of one reactant with an amino group of the other reactant to form a linking amide bond. The amide bond formation may be performed by reacting the starting materials in the presence of a coupling agent or by converting the carboxyl functionality into an active form such as an active ester, mixed anhydride or a carboxyl acid chloride or bromide. General descriptions of such
20 coupling reactions and the reagents used therein can be found in general textbooks on peptide chemistry, for example, M. Bodanszky, "Peptide Chemistry", 2nd rev. ed., Springer-Verlag, Berlin, Germany, (1993).

25 Examples of coupling reactions with amide bond formation include the azide method, mixed carbonic-carboxylic acid anhydride (isobutyl chloroformate) method, the carbodiimide (dicyclohexylcarbodiimide, diisopropylcarbodiimide, or water-soluble carbodiimide such as *N*-ethyl-*N'*-[(3-dimethylamino)propyl]carbodiimide) method, the active ester method (e.g. *p*-nitrophenyl, *p*-chlorophenyl, trichlorophenyl, pentachlorophenyl, pentafluorophenyl, *N*-hydroxysuccinic imido and the like esters), the
30 Woodward reagent K-method, the 1,1-carbonyldiimidazole (CDI or *N,N'*-carbonyldiimidazole) method, the phosphorus reagents or oxidation-reduction methods. Some of these methods can be enhanced by adding suitable catalysts, e.g. in the carbodiimide method by adding 1-hydroxybenzotriazole, DBU (1,8-diazabicyclo[5.4.0]undec-7-ene), or 4-DMAP. Further coupling agents are (benzotriazol-1-yloxy)tris-(dimethylamino)
35 phosphonium hexafluorophosphate, either by itself or in the presence of 1-hydroxybenzotriazole or 4-DMAP; or 2-(1*H*-benzotriazol-1-yl)-*N,N,N',N'*-tetra-methyluronium tetrafluoroborate, or O-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium

hexafluorophosphate. These coupling reactions can be performed in either solution (liquid phase) or solid phase.

A preferred amide bond formation is performed employing N-ethyloxycarbonyl-
5 2-ethyloxy-1,2-dihydroquinoline (EEDQ) or N-isobutyloxy-carbonyl-2-isobutyloxy-
1,2-dihydroquinoline (IIDQ). Unlike the classical anhydride procedure, EEDQ and
IIDQ do not require base nor low reaction temperatures. Typically, the procedure
involves reacting equimolar amounts of the carboxyl and amine components in an
organic solvent (a wide variety of solvents can be used). Then EEDQ or IIDQ is added
10 in excess and the mixture is allowed to stir at room temperature.

The coupling reactions preferably are conducted in an inert solvent, such as
halogenated hydrocarbons, e.g. dichloromethane, chloroform, dipolar aprotic solvents
such as acetonitrile, dimethylformamide, dimethylacetamide, DMSO, HMPT, ethers
15 such as tetrahydrofuran (THF).

In many instances the coupling reactions are done in the presence of a suitable base
such as a tertiary amine, e.g. triethylamine, diisopropylethylamine (DIPEA), *N*-methyl-
morpholine, *N*-methylpyrrolidine, 4-DMAP or 1,8-diazabicyclo[5.4.0]undec-7-ene
20 (DBU). The reaction temperature may range between 0 °C and 50 °C and the reaction
time may range between 15 min and 24 h.

The functional groups in the building blocks that are linked together may be protected
to avoid formation of undesired bonds. Appropriate protecting groups that can be used
25 are listed for example in Greene, "Protective Groups in Organic Chemistry", John
Wiley & Sons, New York (1999) and "The Peptides: Analysis, Synthesis, Biology",
Vol. 3, Academic Press, New York (1987).

Carboxyl groups can be protected as an ester that can be cleaved off to give the
30 carboxylic acid. Protecting groups that can be used include 1) alkyl esters such as
methyl, trimethylsilyl and *tert*-butyl; 2) arylalkyl esters such as benzyl and substituted
benzyl; or 3) esters that can be cleaved by a mild base or mild reductive means such as
trichloroethyl and phenacyl esters.

35 Amino groups can be protected by a variety of N-protecting groups, such as:
1) acyl groups such as formyl, trifluoroacetyl, phthalyl, and *p*-toluenesulfonyl;
2) aromatic carbamate groups such as benzyloxycarbonyl (Cbz or Z) and
substituted benzyloxycarbonyls, and 9-fluorenylmethyloxycarbonyl (Fmoc);

- 3) aliphatic carbamate groups such as *tert*-butyloxycarbonyl (Boc),
ethoxycarbonyl, diisopropylmethoxy-carbonyl, and allyloxycarbonyl;
 - 4) cyclic alkyl carbamate groups such as cyclopentyloxycarbonyl and
adamantyloxycarbonyl;
 - 5) 5) alkyl groups such as triphenylmethyl, benzyl or substituted benzyl such as
4-methoxybenzyl;
 - 6) trialkylsilyl such as trimethylsilyl or *t*.Bu dimethylsilyl; and
 - 7) thiol containing groups such as phenylthiocarbonyl and dithiasuccinoyl.
- Interesting amino protecting groups are Boc and Fmoc.

10

Preferably the amino protecting group is cleaved off prior to the next coupling step. Removal of N-protecting groups can be done following art-known procedures. When the Boc group is used, the methods of choice are trifluoroacetic acid, neat or in dichloromethane, or HCl in dioxane or in ethyl acetate. The resulting ammonium salt is then neutralized either prior to the coupling or in situ with basic solutions such as aqueous buffers, or tertiary amines in dichloromethane or acetonitrile or dimethylformamide. When the Fmoc group is used, the reagents of choice are piperidine or substituted piperidine in dimethylformamide, but any secondary amine can be used. The deprotection is carried out at a temperature between 0 °C and room temperature, usually around 15-25 °C, or 20-22 °C.

20

Other functional groups that can interfere in the coupling reactions of the building blocks may also be protected. For example hydroxyl groups may be protected as benzyl or substituted benzyl ethers, e.g. 4-methoxybenzyl ether, benzoyl or substituted benzoyl esters, e.g. 4-nitrobenzoyl ester, or with trialkylsilyl groups (e.g. trimethylsilyl or *tert*-butyldimethylsilyl).

25

Further amino groups may be protected by protecting groups that can be cleaved off selectively. For example, when Boc is used as the α -amino protecting group, the following side chain protecting groups are suitable: *p*-toluenesulfonyl (tosyl) moieties can be used to protect further amino groups; benzyl (Bn) ethers can be used to protect hydroxy groups; and benzyl esters can be used to protect further carboxyl groups. Or when Fmoc is chosen for the α -amino protection, usually *tert*-butyl based protecting groups are acceptable. For instance, Boc can be used for further amino groups; *tert*-butyl ethers for hydroxyl groups; and *tert*-butyl esters for further carboxyl groups.


30

35

Any of the protecting groups may be removed at any stage of the synthesis procedure but preferably, the protecting groups of any of the functionalities not involved in the

reaction steps are removed after completion of the build-up of the macrocycle. Removal of the protecting groups can be done in whatever manner is dictated by the choice of protecting groups, which manners are well known to those skilled in the art.


- 5 The intermediates of formula (1a) wherein X is N, said intermediates being represented by formula (1a-1), may be prepared using an urea forming reaction, starting from intermediates (5a) which are reacted with an alkenamine (5b) in the presence of a carbonyl introducing agent as outlined in the following reaction scheme.



- 10 Carbonyl (CO) introducing agents include phosgene, or phosgene derivatives such as carbonyl diimidazole (CDI), and the like. In one embodiment (5a) is reacted with the CO introducing agent in the presence of a suitable base and a solvent, which can be the bases and solvents used in the amide forming reactions as described above. Thereafter, the amine (5b) is added thereby obtaining intermediates (1a-1) as in the above scheme.
- 15 In a particular embodiment, the base is a hydrogencarbonate, e.g. NaHCO_3 , or a tertiary amine such as triethylamine and the like, and the solvent is an ether or halogenated hydrocarbon, e.g. THF, CH_2Cl_2 , CHCl_3 , and the like. An alternative route using similar reaction conditions involves first reacting the CO introducing agent with the amine (5b) and then reacting the thus formed intermediate with (5a).


20

The intermediates (1a-1) can alternatively be prepared as follows:




- PG¹ is an O-protecting group, which can be any of the groups mentioned herein and in particular is a benzoyl or substituted benzoyl group such as 4-nitrobenzoyl. In the latter instance this group can be removed by reaction with an alkali metal hydroxide (LiOH, NaOH, KOH), in particular where PG¹ is 4-nitrobenzoyl, with LiOH, in an aqueous medium comprising water and a water-soluble organic solvent such as an alkanol (methanol, ethanol) and THF.
- Intermediates (6a) are reacted with (5b) in the presence of a carbonyl introducing agent, similar as described above, and this reaction yields intermediates (6c). These are deprotected, in particular using the reaction conditions mentioned above. The resulting alcohol (6d) is reacted with intermediates (4b) in a carbamate forming reaction, as described above for the reaction of (4a) with (4b), and this reaction results in intermediates (1a-1).


The intermediates of formula (1a) wherein X is C, said intermediates being represented by formula (1a-2), may be prepared by an amide forming reaction starting from intermediates (7a) which are reacted with an alkenamine (5b) as shown in the following reaction scheme, using reaction conditions for preparing amides such as those described above.



The intermediates (1a-1) can alternatively be prepared as follows:




- 5 PG¹ is an O-protecting group as described above. The same reaction conditions as described above may be used: amide formation as described above, removal of PG¹ as in the description of the protecting groups and introduction of R⁸ as in the reactions of (4a) with the amine s (4b).
- 10 The intermediates of formula (2a) may be prepared by first cyclizing an open amide (9a) to a macrocyclic ester (9b), which in turn is converted to an intermediate (2a) as follows:



PG² is a carboxyl protecting group, e.g. one of the carboxyl protecting groups mentioned above, in particular a C₁₋₄alkyl or benzyl ester, e.g. a methyl, ethyl or t.butyl ester. The reaction of (9a) to (9b) is a metathesis reaction and is conducted as described above. Removal of PG² as described above, yields intermediates (2a). Where PG¹ is a C₁₋₄alkyl ester, it is removed by alkaline hydrolysis, e.g. with NaOH or preferably LiOH, in an aqueous solvent, e.g. a C₁₋₄alkanol/water mixture, such as methanol/water or ethanol/water. A benzyl group can be removed by catalytic hydrogenation.

10

In an alternative synthesis, intermediates (2a) can be prepared as follows:



The PG¹ group is selected such that it is selectively cleavable towards PG². PG² may be e.g. methyl or ethyl esters, which can be removed by treatment with an alkali metal


hydroxide in an aqueous medium, in which case PG^1 e.g. is t.butyl or benzyl. Or alternatively, PG^2 may be t.butyl esters removable under weakly acidic conditions or PG^1 may be benzyl esters removable with strong acid or by catalytic hydrogenation, in the latter two cases PG^1 e.g. is a benzoic ester such as a 4-nitrobenzoic ester.

5

First, intermediates (10a) are cyclized to the macrocyclic esters (10b), the latter are deprotected by removal of the PG^1 group to intermediates (10c), which are reacted with amine s (4b), followed by removal of the carboxyl protecting group PG^2 . The cyclization, deprotection of PG^1 and PG^2 , and the coupling with (4b) are as described above.


10

The R^1 groups can be introduced at any stage of the synthesis, either as the last step as described above, or earlier, before the macrocycle formation, as illustrated in the following scheme:



15

In the above scheme, R^2 , R^6 , R^7 , R^8 , X and PG^2 are as defined above and L^1 is a P3 group



wherein n and R^3 are as defined above and where X is N, L^1 may also be a nitrogen-protecting group (PG, as defined above) and where X is C, L^1 may also be a group


20

–COOPG^{2a}, wherein the group PG^{2a} is a carboxyl protecting group similar as PG², but wherein PG^{2a} is selectively cleavable towards PG². In one embodiment PG^{2a} is t.butyl and PG² is methyl or ethyl.

- 5 The intermediates (11c) and (11d) wherein L¹ represents a group (b) correspond to the intermediates (1a) and may be processed further as specified above.

Coupling of P1 and P2 building blocks


- The P1 and P2 building blocks are linked using an amide forming reaction following the procedures described above. The P1 building block may have a carboxyl protecting group PG² (as in (12b)) or may already be linked to P1' group (as in (12c)). L² is a N-protecting group (PG), or a group (b), as specified above. L³ is hydroxy, -OPG¹ or a group -O-R⁸ as specified above. Where in any of the following reaction schemes L³ is hydroxy, prior to each reaction step, it may be protected as a group -OPG¹ and, if
- 10 described, subsequently deprotected back to a free hydroxy function. Similarly as
- 15 described above, the hydroxy function may be converted to a group -O-R⁸.



- In the procedure of the above scheme, a cyclopropyl amino acid (12b) or (12c) is coupled to the acid function of the P2 building block (12a) with the formation of an amide linkage, following the procedures described above. Intermediates (12d) or (12e) are obtained. Where in the latter L² is a group (b), the resulting products are P3-P2-P1 sequences encompassing some of the intermediates (11c) or (11d) in the previous reaction scheme. Removal of the acid protecting group in (12d), using the appropriate conditions for the protecting group used, followed by coupling with an amine
- 20


H₂N-SO₂R⁷ (2b) or with HOR⁶ (2c) as described above, again yields the intermediates (12e), wherein -COR¹ are amide or ester groups. Where L² is a N-protecting group, it can be removed yielding intermediates (5a) or (6a). In one embodiment, PG in this reaction is a BOC group and PG² is methyl or ethyl. Where additionally L³ is hydroxy, the starting material (12a) is Boc-L-hydroxyproline. In a particular embodiment, PG is BOC, PG² is methyl or ethyl and L³ is -O-R⁸.

In one embodiment, L² is a group (b) and these reactions involve coupling P1 to P2-P3, which results in the intermediates (1a-1) or (1a) mentioned above. In another embodiment, L² is a N-protecting group PG, which is as specified above, and the coupling reaction results in intermediates (12d-1) or (12e-1), from which the group PG can be removed, using reaction conditions mentioned above, obtaining intermediates (12-f) or respectively (12g), which encompass intermediates (5a) and (6a) as specified above:




In one embodiment, the group L³ in the above schemes represents a group -O-PG¹ which can be introduced on a starting material (12a) wherein L³ is hydroxy. In this instance PG¹ is chosen such that it is selectively cleavable towards group L² being PG.


In a similar way, P2 building blocks wherein X is C, which are cyclopentane or cyclopentene derivatives, can be linked to P1 building blocks as outlined in the following scheme wherein R¹, R², L³, PG² and PG^{2a} are carboxyl protecting groups. PG^{2a} typically is chosen such that it is selectively cleavable towards group PG². Removal of the PG^{2a} group in (13c) yields intermediates (7a) or (8a), which can be reacted with (5b) as described above.



In one particular embodiment, where X is C, R^2 is H, and where X and the carbon bearing R^2 are linked by a single bond (P2 being a cyclopentane moiety), PG^{2a} and L^3 taken together form a bond and the P2 building block is represented by formula:



Bicyclic acid (14a) is reacted with (12b) or (12c) similar as described above to (14b) and (14c) respectively, wherein the lactone is opened to give intermediates (14c) and (14e). The lactone can be opened using ester hydrolysis procedures, for example using the reaction conditions described above for the alkaline removal of a PG^1 group in (9b),
 10 in particular using basic conditions such as an alkali metal hydroxide, e.g. NaOH, KOH, in particular LiOH.



Intermediates (14c) and (14e) can be processed further as described hereinafter.


Synthesis of P2 building blocks

- 5 The P2 building blocks contain either a pyrrolidine, a cyclopentane, or a cyclopentene moiety substituted with a group $-O-R^8$.

P2 building blocks containing a pyrrolidine moiety can be derived from commercially available hydroxy proline.

10


The preparation of P2 building blocks that contain a cyclopentane ring may be performed as shown in the scheme below.




The bicyclic acid (17b) can be prepared, for example, from 3,4-bis(methoxycarbonyl)-cyclopentanone (17a), as described by Rosenquist et al. in Acta Chem. Scand. 46 (1992) 1127-1129. A first step in this procedure involves the reduction of the keto group with a reducing agent like sodium borohydride in a solvent such as methanol, followed by hydrolysis of the esters and finally ring closure to the bicyclic lactone (17b) using lactone forming procedures, in particular by using acetic anhydride in the presence of a weak base such as pyridine. The carboxylic acid functionality in (17b) can then be protected by introducing an appropriate carboxyl protecting group, such as a group PG², which is as specified above, thus providing bicyclic ester (17c). The group PG² in particular is acid-labile such as a t.butyl group and is introduced e.g. by treatment with isobutene in the presence of a Lewis acid or with di-*tert*-butyl dicarbonate in the presence of a base such as a tertiary amine like dimethylamino-pyridine or triethylamine in a solvent like dichloromethane. Lactone opening of (17c) using reaction conditions described above, in particular with lithium hydroxide, yields the acid (17d), which can be used further in coupling reactions with P1 building blocks. The free acid in (17d) may also be protected, preferably with an acid protecting group PG^{2a} that is selectively cleavable towards PG², and the hydroxy function may be converted to a group -OPG¹ or to a group -O-R⁸. The products obtained upon removal of the group PG² are intermediates (17g) and (17i) which correspond to intermediates (13a) or (16a) specified above.

Intermediates with specific stereochemistry may be prepared by resolving the intermediates in the above reaction sequence. For example, (17b) may be resolved following art-known procedures, e.g. by salt formation with an optically active base or by chiral chromatography, and the resulting stereoisomers may be processed further as described above. The OH and COOH groups in (17d) are in cis position. Trans analogs can be prepared by inverting the stereochemistry at the carbon bearing the OH function by using specific reagents in the reactions introducing OPG¹ or O-R⁸ that invert the stereochemistry, such as, e.g. by applying a Mitsunobu reaction.

In one embodiment, the intermediates (17d) are coupled to P1 blocks (12b) or (12c), which coupling reactions correspond to the coupling of (13a) or (16a) with the same P1 blocks, using the same conditions. Subsequent introduction of a -O-R⁸-substituent as described above followed by removal of the acid protection group PG² yields intermediates (8a-1), which are a subclass of the intermediates (7a), or part of the intermediates (16a). The reaction products of the PG² removal can be further coupled to a P3 building block. In one embodiment PG² in (17d) is t.butyl which can be removed under acidic conditions, e.g. with trifluoroacetic acid.




An unsaturated P2 building block, i.e. a cyclopentene ring, may be prepared as illustrated in the scheme below.



- 5 A bromination-elimination reaction of 3,4-bis(methoxycarbonyl)cyclopentanone (17a) as described by Dolby et al. in J. Org. Chem. 36 (1971) 1277-1285 followed by reduction of the keto functionality with a reducing agent like sodium borohydride provides the cyclopentenol (19a). Selective ester hydrolysis using for example lithium hydroxide in a solvent like a mixture of dioxane and water, provides the hydroxy substituted monoester cyclopentenol (19b).
- 10

An unsaturated P2 building block wherein R² can also be other than hydrogen, may be prepared as shown in the scheme below.




- 15 Oxidation of commercially available 3-methyl-3-buten-1-ol (20a), in particular by an oxidizing agent like pyridinium chlorochromate, yields (20b), which is converted to the

corresponding methyl ester, e.g. by treatment with acetyl chloride in methanol, followed by a bromination reaction with bromine yielding the α -bromo ester (20c). The latter can then be condensed with the alkenyl ester (20e), obtained from (20d) by an ester forming reaction. The ester in (20e) preferably is a t.butyl ester which can be prepared from the corresponding commercially available acid (20d), e.g. by treatment with di-*tert*-butyl dicarbonate in the presence of a base like dimethylaminopyridine. Intermediate (20e) is treated with a base such as lithium diisopropyl amide in a solvent like tetrahydrofuran, and reacted with (20c) to give the alkenyl diester (20f). Cyclisation of (20f) by an olefin metathesis reaction, performed as described above, provides cyclopentene derivative (20g). Stereoselective epoxidation of (20g) can be carried out using the Jacobsen asymmetric epoxidation method to obtain epoxide (20h). Finally, an epoxide opening reaction under basic conditions, e.g. by addition of a base, in particular DBN (1,5-diazabicyclo-[4.3.0]non-5-ene), yields the alcohol (20i). Optionally, the double bond in intermediate (20i) can be reduced, for example by catalytic hydrogenation using a catalyst like palladium on carbon, yielding the corresponding cyclopentane compound. The t.butyl ester may be removed to the corresponding acid, which subsequently is coupled to a P1 building block.

The $-O-R^8$ group can be introduced on the pyrrolidine, cyclopentane or cyclopentene rings at any convenient stage of the synthesis of the compounds according to the present invention. One approach is to first introduce the $-O-R^8$ group to the said rings and subsequently add the other desired building blocks, i.e. P1 (optionally with the P1' tail) and P3, followed by the macrocycle formation. Another approach is to couple the building blocks P2, bearing no $-O-R^8$ substituent, with each P1 and P3, and to add the $-O-R^8$ group either before or after the macrocycle formation. In the latter procedure, the P2 moieties have a hydroxy group, which may be protected by a hydroxy protecting group PG¹.


R^8 groups can be introduced on building blocks P2 by reacting hydroxy substituted intermediates (21a) or (21b) with intermediates (4b) similar as described above for the synthesis of (I) starting from (4a). These reactions are represented in the schemes below, wherein L^2 is as specified above and L^5 and L^{5a} independently from one another, represent hydroxy, a carboxyl protecting group $-OPG^2$ or $-OPG^{2a}$, or L^5 may also represent a P1 group such as a group (d) or (e) as specified above, or L^{5a} may also represent a P3 group such as a group (b) as specified above. The groups PG² and PG^{2a} are as specified above. Where the groups L^5 and L^{5a} are PG² or PG^{2a}, they are chosen such that each group is selectively cleavable towards the other. For example, one of L^5 and L^{5a} may be a methyl or ethyl group and the other a benzyl or t.butyl group.

In one embodiment in (21a), L^2 is PG and L^5 is $-OPG^2$, or in (21d), L^{5a} is $-OPG^2$ and L^5 is $-OPG^2$ and the PG^2 groups are removed as described above.




(21a)

(21b)




(21b-1)

(21c)



(21d)

(21e)



(21e-1)

(21f)

In another embodiment the group L^2 is BOC, L^5 is hydroxy and the starting material (21a) is commercially available BOC-hydroxyproline, or any other stereoisomeric form thereof, e.g. BOC-L-hydroxyproline, in particular the trans isomer of the latter. Where L^5 in (21b) is a carboxyl-protecting group, it may be removed following procedures described above to (21c). In still another embodiment PG in (21b-1) is Boc and PG^2 is a lower alkyl ester, in particular a methyl or ethyl ester. Hydrolysis of the latter ester to the acid can be done by standard procedures, e.g. acid hydrolysis with hydrochloric acid in methanol or with an alkali metal hydroxide such as NaOH, in particular with

LiOH. In another embodiment, hydroxy substituted cyclopentane or cyclopentene analogs (21d) are converted to (21e), which, where L^5 and L^{5a} are $-OPG^2$ or $-OPG^{2a}$, may be converted to the corresponding acids (21f) by removal of the group PG^2 . Removal of PG^{2a} in (21e-1) leads to similar intermediates.

5


Intermediates (4b), which are amino derivatives, are known compounds or can be easily prepared using art-known procedures.

Synthesis of P1 building blocks

- 10 The cyclopropane amino acid used in the preparation of the P1 fragment is commercially available or can be prepared using art-known procedures.

In particular the amino-vinyl-cyclopropyl ethyl ester (12b) may be obtained according to the procedure described in WO 00/09543 or as illustrated in the following scheme, wherein PG^2 is a carboxyl protecting group as specified above:


15



- Treatment of commercially available or easily obtainable imine (31a) with 1,4-dihalo-butene in presence of a base produces (31b), which after hydrolysis yields cyclopropyl amino acid (12b), having the allyl substituent *syn* to the carboxyl group. Resolution of the enantiomeric mixture (12b) results in (12b-1). The resolution is performed using art-known procedures such as enzymatic separation; crystallization with a chiral acid; or chemical derivatization; or by chiral column chromatography. Intermediates (12b) or (12b-1) may be coupled to the appropriate P2 derivatives as described above.

- 25 P1 building blocks for the preparation of compounds according to general formula (I) wherein R^1 is $-OR^6$ or $-NH-SO_2R^7$ can be prepared by reacting amino acids (32a) with the appropriate alcohol or amine respectively under standard conditions for ester or amide formation. Cyclopropyl amino acids (32a) are prepared by introducing a N-

protecting group PG, and removal of PG² and the amino acids (32a) are converted to the amides (12c-1) or esters (12c-2), which are subgroups of the intermediates (12c), as outlined in the following reaction scheme, wherein PG is as specified above.



- 5 The reaction of (32a) with amine (2b) is an amide forming procedure. The similar reaction with (2c) is an ester forming reaction. Both can be performed following the procedures described above. This reaction yields intermediates (32b) or (32c) from which the amino protecting group is removed by standard methods such as those described above. This in turn results in the desired intermediate (12c-1). Starting materials (32a) may be prepared from the above mentioned intermediates (12b) by first introducing a N-protecting group PG and subsequent removal of the group PG².
- 10

In one embodiment the reaction of (32a) with (2b) is done by treatment of the amino acid with a coupling agent, for example N,N'-carbonyl-diimidazole (CDI) or the like, in a solvent like THF followed by reaction with (2b) in the presence of a base such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). Alternatively the amino acid can be treated with (2b) in the presence of a base like diisopropylethylamine followed by treatment with a coupling agent such as benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate (commercially available as PyBOP®) to effect the introduction of the sulfonamide group.


20

Intermediates (12c-1) or (12c-2) in turn may be coupled to the appropriate proline, cyclopentane or cyclopentene derivatives as described above.

25 Synthesis of the P3 building blocks

The P3 building blocks are available commercially or can be prepared according to methodologies known to the skilled in the art. One of these methodologies is shown in


the scheme below and uses monoacylated amines, such as trifluoroacetamide or a Boc-protected amine.



In the above scheme, R together with the CO group forms a N-protecting group, in particular R is *t*-butoxy, trifluoromethyl; R^3 and n are as defined above and LG is a leaving group, in particular halogen, e.g. chloro or bromo.

The monoacylated amines (33a) are treated with a strong base such as sodium hydride and are subsequently reacted with a reagent LG- C_{5-8} alkenyl (33b), in particular halo C_{5-8} alkenyl, to form the corresponding protected amines (33c). Deprotection of (33c) affords (5b), which are building blocks P3. Deprotection will depend on the functional group R, thus if R is *t*-butoxy, deprotection of the corresponding Boc-protected amine can be accomplished with an acidic treatment, e.g. trifluoroacetic acid. Alternatively, when R is for instance trifluoromethyl, removal of the R group is accomplished with a base, e.g. sodium hydroxide.

The following scheme illustrates yet another method for preparing a P3 building block, namely a Gabriel synthesis of primary C_{5-8} alkenylamines, which can be carried out by the treatment of a phthalimide (34a) with a base, such as NaOH or KOH, and with (33b), which is as specified above, followed by hydrolysis of the intermediate N-alkenyl imide to generate a primary C_{5-8} alkenylamine (5b-1).



In the above scheme, n is as defined above.

Compounds of formula (I) may be converted into each other following art-known functional group transformation reactions. For example, amino groups may be N-alkylated, nitro groups reduced to amino groups, a halo atom may be exchanged for another halo.

The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate
5 inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarbo-
peroxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzene-
carboperoxoic acid, peroalkanoic acids, e.g. peroacetic acid, alkylhydroperoxides,
10 e.g. *tert*-butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained
15 by the application of art-known procedures. Diastereomers may be separated by physical methods such as selective crystallization and chromatographic techniques, e.g., counter-current distribution, liquid chromatography and the like.

The compounds of formula (I) may be obtained as racemic mixtures of enantiomers
20 which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I), which are sufficiently basic or acidic may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid, respectively chiral base. Said diastereomeric salt forms are subsequently
separated, for example, by selective or fractional crystallization and the enantiomers are
25 liberated therefrom by alkali or acid. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase. Said pure
stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the
30 reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound may be synthesized by stereospecific methods of preparation. These methods may advantageously employ enantiomerically pure starting materials.

In a further aspect, the present invention concerns a pharmaceutical composition
35 comprising a therapeutically effective amount of a compound of formula (I) as specified herein, or a compound of any of the subgroups of compounds of formula (I) as specified herein, and a pharmaceutically acceptable carrier. A therapeutically effective amount in this context is an amount sufficient to prophylactically act against,

to stabilize or to reduce viral infection, and in particular HCV viral infection, in infected subjects or subjects being at risk of being infected. In still a further aspect, this invention relates to a process of preparing a pharmaceutical composition as specified herein, which comprises intimately mixing a pharmaceutically acceptable carrier with a therapeutically effective amount of a compound of formula (I), as specified herein, or of a compound of any of the subgroups of compounds of formula (I) as specified herein.

Therefore, the compounds of the present invention or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form or metal complex, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin.

The compounds of the present invention may also be administered via oral inhalation or insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry powder, a solution being preferred. Any system developed for the delivery of solutions, suspensions or dry powders via oral inhalation or insufflation are suitable for the administration of the present compounds.

Thus, the present invention also provides a pharmaceutical composition adapted for administration by inhalation or insufflation through the mouth comprising a compound of formula (I) and a pharmaceutically acceptable carrier. Preferably, the compounds of the present invention are administered via inhalation of a solution in nebulized or aerosolized doses.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, suppositories, powder packets, wafers, injectable solutions or suspensions and the like, and segregated multiples thereof.

The compounds of formula (I) show antiviral properties. Viral infections and their associated diseases treatable using the compounds and methods of the present invention include those infections brought on by HCV and other pathogenic flaviviruses such as Yellow fever, Dengue fever (types 1-4), St. Louis encephalitis, Japanese encephalitis, Murray valley encephalitis, West Nile virus and Kunjin virus. The diseases associated with HCV include progressive liver fibrosis, inflammation and necrosis leading to cirrhosis, end-stage liver disease, and HCC; and for the other pathogenic flaviviruses the diseases include yellow fever, dengue fever, hemorrhagic fever and encephalitis. A number of the compounds of this invention moreover are active against mutated strains of HCV. Additionally, many of the compounds of this invention show a favorable pharmacokinetic profile and have attractive properties in terms of bioavailability, including an acceptable half-life, AUC (area under the curve) and peak values and lacking unfavorable phenomena such as insufficient quick onset and tissue retention.

The *in vitro* antiviral activity against HCV of the compounds of formula (I) was tested in a cellular HCV replicon system based on Lohmann et al. (1999) Science 285:110-113, with the further modifications described by Krieger et al. (2001) Journal of Virology 75: 4614-4624, which is further exemplified in the examples section. This model, while not a complete infection model for HCV, is widely accepted as the most robust and efficient model of autonomous HCV RNA replication currently available. Compounds exhibiting anti-HCV activity in this cellular model are considered as candidates for further development in the treatment of HCV infections in mammals. It will be appreciated that it is important to distinguish between compounds that specifically interfere with HCV functions from those that exert cytotoxic or cytostatic effects in the HCV replicon model, and as a consequence cause a decrease in HCV RNA or linked reporter enzyme concentration. Assays are known in the field for the evaluation of cellular cytotoxicity based for example on the activity of mitochondrial enzymes using fluorogenic redox dyes such as resazurin. Furthermore, cellular counter screens exist for the evaluation of non-selective inhibition of linked reporter gene activity, such as firefly luciferase. Appropriate cell types can be equipped by stable transfection with a luciferase reporter gene whose expression is dependent on a constitutively active gene promoter, and such cells can be used as a counter-screen to eliminate non-selective inhibitors.

Due to their antiviral properties, particularly their anti-HCV properties, the compounds of formula (I) or any subgroup thereof, their prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms, are useful in the treatment of individuals experiencing a viral infection, particularly a HCV infection, and for the prophylaxis of these infections. In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected with viruses, in particular flaviviruses such as HCV.

The compounds of the present invention or any subgroup thereof may therefore be used as medicines. Said use as a medicine or method of treatment comprises the systemic administration to viral infected subjects or to subjects susceptible to viral infections of an amount effective to combat the conditions associated with the viral infection, in particular the HCV infection.

The present invention also relates to the use of the present compounds or any subgroup thereof in the manufacture of a medicament for the treatment or the prevention of viral infections, particularly HCV infection.

The present invention furthermore relates to a method of treating a warm-blooded animal infected by a virus, or being at risk of infection by a virus, in particular by HCV, said method comprising the administration of an anti-virally effective amount of a compound of formula (I), as specified herein, or of a compound of any of the
5 subgroups of compounds of formula (I), as specified herein.

Also, the combination of previously known anti-HCV compound, such as, for instance, interferon- α (IFN- α), pegylated interferon- α and/or ribavirin, and a compound of formula (I) can be used as a medicine in a combination therapy. The term
10 "combination therapy" relates to a product containing mandatory (a) a compound of formula (I), and (b) optionally another anti-HCV compound, as a combined preparation for simultaneous, separate or sequential use in treatment of HCV infections, in particular, in the treatment of infections with HCV.

15 Anti-HCV compounds encompass agents selected from an HCV polymerase inhibitor, an HCV protease inhibitor, an inhibitor of another target in the HCV life cycle, and immunomodulatory agent, an antiviral agent, and combinations thereof.

HCV polymerase inhibitors include, but are not limited to, NM283 (valopicitabine),
20 R803, JTK-109, JTK-003, HCV-371, HCV-086, HCV-796 and R-1479.

Inhibitors of HCV proteases (NS2-NS3 inhibitors and NS3-NS4A inhibitors) include, but are not limited to, the compounds of WO02/18369 (see, e.g., page 273, lines 9-22 and page 274, line 4 to page 276, line 11); BILN-2061, VX-950, GS-9132 (ACH-806),
25 SCH-503034, and SCH-6. Further agents that can be used are those disclosed in WO98/17679, WO00/056331 (Vertex); WO 98/22496 (Roche); WO 99/07734, (Boehringer Ingelheim), WO 2005/073216, WO 2005073195 (Medivir) and structurally similar agents.

30 Inhibitors of other targets in the HCV life cycle, including NS3 helicase; metallo-protease inhibitors; antisense oligonucleotide inhibitors, such as ISIS-14803, AVI-4065 and the like; siRNA's such as SIRPLEX-140-N and the like; vector-encoded short hairpin RNA (shRNA); DNazymes; HCV specific ribozymes such as heptazyme, RPI.13919 and the like; entry inhibitors such as HepeX-C, HuMax-HepC and the like;
35 alpha glucosidase inhibitors such as celgosivir, UT-231B and the like; KPE-02003002; and BIVN 401.

Immunomodulatory agents include, but are not limited to; natural and recombinant interferon isoform compounds, including α -interferon, β -interferon, γ -interferon, ω -interferon and the like, such as Intron A®, Roferon-A®, Canferon-A300®, Advaferon®, Infergen®, Humoferon®, Sumiferon MP®, Alfaferone®, IFN-beta®,
5 Feron® and the like; polyethylene glycol derivatized (pegylated) interferon compounds, such as PEG interferon- α -2a (Pegasys®), PEG interferon- α -2b (PEG-Intron®), pegylated IFN- α -con1 and the like; long acting formulations and derivatizations of interferon compounds such as the albumin-fused interferon albuferon α and the like; compounds that stimulate the synthesis of interferon in cells, such as
10 resiquimod and the like; interleukins; compounds that enhance the development of type 1 helper T cell response, such as SCV-07 and the like; TOLL-like receptor agonists such as CpG-10101 (actilon), isatoribine and the like; thymosin α -1; ANA-245; ANA-246; histamine dihydrochloride; propagermanium; tetrachlorodecaoxide; ampligen; IMP-321; KRN-7000; antibodies, such as civacir, XTL-6865 and the like; and
15 prophylactic and therapeutic vaccines such as InnoVac C, HCV E1E2/MF59 and the like.

Other antiviral agents include, but are not limited to, ribavirin, amantadine, viramidine, nitazoxanide; telbivudine; NOV-205; taribavirin; inhibitors of internal ribosome entry;
20 broad-spectrum viral inhibitors, such as IMPDH inhibitors (e.g., compounds of US5,807,876, US6,498,178, US6,344,465, US6,054,472, WO97/40028, WO98/40381, WO00/56331, and mycophenolic acid and derivatives thereof, and including, but not limited to VX-950, merimepodib (VX-497), VX-148, and/or VX-944); or combinations of any of the above.

25 Thus, to combat or treat HCV infections, the compounds of formula (I) may be co-administered in combination with for instance, interferon- α (IFN- α), pegylated interferon- α and/or ribavirin, as well as therapeutics based on antibodies targeted against HCV epitopes, small interfering RNA (si RNA), ribozymes, DNazymes,
30 antisense RNA, small molecule antagonists of for instance NS3 protease, NS3 helicase and NS5B polymerase.

Accordingly, the present invention relates to the use of a compound of formula (I) or any subgroup thereof as defined above for the manufacture of a medicament useful for
35 inhibiting HCV activity in a mammal infected with HCV viruses, wherein said medicament is used in a combination therapy, said combination therapy preferably comprising a compound of formula (I) and another HCV inhibitory compound, e.g. (pegylated) IFN- α and/or ribavirin.


In still another aspect there are provided combinations of a compound of formula (I) as specified herein and an anti-HIV compound. The latter preferably are those HIV inhibitors that have a positive effect on drug metabolism and/or pharmacokinetics that improve bioavailability. An example of such an HIV inhibitor is ritonavir.

As such, the present invention further provides a combination comprising (a) an HCV NS3/4a protease inhibitor of formula (I) or a pharmaceutically acceptable salt thereof; and (b) ritonavir or a pharmaceutically acceptable salt thereof.

10

The compound ritonavir, and pharmaceutically acceptable salts thereof, and methods for its preparation are described in WO 94/14436. For preferred dosage forms of ritonavir, see US 6,037, 157, and the documents cited therein: US 5,484, 801, US 08/402,690, and WO 95/07696 and WO 95/09614. Ritonavir has the following formula:

15



In a further embodiment, the combination comprising (a) an HCV NS3/4a protease inhibitor of formula (I) or a pharmaceutically acceptable salt thereof; and (b) ritonavir or a pharmaceutically acceptable salt thereof; further comprises an additional anti-HCV compound selected from the compounds as described herein.

20

In one embodiment of the present invention there is provided a process for preparing a combination as described herein, comprising the step of combining an HCV NS3/4a protease inhibitor of formula (I) or a pharmaceutically acceptable salt thereof, and ritonavir or a pharmaceutically acceptable salt thereof. An alternative embodiment of this invention provides a process wherein the combination comprises one or more additional agent as described herein.

25

The combinations of the present invention may be used as medicaments. Said use as a medicine or method of treatment comprises the systemic administration to HCV-infected subjects of an amount effective to combat the conditions associated with HCV

30

and other pathogenic flavi- and pestiviruses. Consequently, the combinations of the present invention can be used in the manufacture of a medicament useful for treating, preventing or combating infection or disease associated with HCV infection in a mammal, in particular for treating conditions associated with HCV and other

5 pathogenic flavi- and pestiviruses.

In one embodiment of the present invention there is provided a pharmaceutical composition comprising a combination according to any one of the embodiments described herein and a pharmaceutically acceptable excipient. In particular, the present invention provides a pharmaceutical composition comprising (a) a therapeutically
10 effective amount of an HCV NS3/4a protease inhibitor of the formula (I) or a pharmaceutically acceptable salt thereof, (b) a therapeutically effective amount of ritonavir or a pharmaceutically acceptable salt thereof, and (c) a pharmaceutically acceptable excipient. Optionally, the pharmaceutical composition further comprises an additional agent selected from an HCV polymerase inhibitor, an HCV protease
15 inhibitor, an inhibitor of another target in the HCV life cycle, and immunomodulatory agent, an antiviral agent, and combinations thereof.

The compositions may be formulated into suitable pharmaceutical dosage forms such as the dosage forms described above. Each of the active ingredients may be formulated
20 separately and the formulations may be co-administered or one formulation containing both and if desired further active ingredients may be provided.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients, as well as any product which results, directly or indirectly,
25 from the combination of the specified ingredients.

In one embodiment the combinations provided herein may also be formulated as a combined preparation for simultaneous, separate or sequential use in HIV therapy. In such a case, the compound of general formula (I) or any subgroup thereof, is
30 formulated in a pharmaceutical composition containing other pharmaceutically acceptable excipients, and ritonavir is formulated separately in a pharmaceutical composition containing other pharmaceutically acceptable excipients. Conveniently, these two separate pharmaceutical compositions can be part of a kit for simultaneous, separate or sequential use.

35

Thus, the individual components of the combination of the present invention can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. The present invention is therefore to be

understood as embracing all such regimes of simultaneous or alternating treatment and the term “administering” is to be interpreted accordingly. In a preferred embodiment, the separate dosage forms are administered about simultaneously.

5 In one embodiment, the combination of the present invention contains an amount of ritonavir, or a pharmaceutically acceptable salt thereof, which is sufficient to clinically improve the bioavailability of the HCV NS3/4a protease inhibitor of formula (I) relative to the bioavailability when said HCV NS3/4a protease inhibitor of formula (I) is administered alone.

10

In another embodiment, the combination of the present invention contains an amount of ritonavir, or a pharmaceutically acceptable salt thereof, which is sufficient to increase at least one of the pharmacokinetic variables of the HCV NS3/4a protease inhibitor of formula (I) selected from $t_{1/2}$, C_{min} , C_{max} , C_{ss} , AUC at 12 hours, or AUC at 24 hours,
15 relative to said at least one pharmacokinetic variable when the HCV NS3/4a protease inhibitor of formula (I) is administered alone.

15

A further embodiment relates to a method for improving the bioavailability of a HCV NS3/4a protease inhibitor comprising administering to an individual in need of such
20 improvement a combination as defined herein, comprising a therapeutically effective amount of each component of said combination.

20

In a further embodiment, the invention relates to the use of ritonavir or a pharmaceutically acceptable salt thereof, as an improver of at least one of the
25 pharmacokinetic variables of a HCV NS3/4a protease inhibitor of formula (I) selected from $t_{1/2}$, C_{min} , C_{max} , C_{ss} , AUC at 12 hours, or AUC at 24 hours; with the proviso that said use is not practised in the human or animal body.

25

The term “individual” as used herein refers to an animal, preferably a mammal, most
30 preferably a human, who has been the object of treatment, observation or experiment.

30

Bioavailability is defined as the fraction of administered dose reaching systemic circulation. $t_{1/2}$ represents the half life or time taken for the plasma concentration to fall to half its original value. C_{ss} is the steady state concentration, i.e. the concentration at
35 which the rate of input of drug equals the rate of elimination. C_{min} is defined as the lowest (minimum) concentration measured during the dosing interval. C_{max} represents the highest (maximum) concentration measured during the dosing interval. AUC is

35

defined as the area under the plasma concentration-time curve for a defined period of time.

5 The combinations of this invention can be administered to humans in dosage ranges specific for each component comprised in said combinations. The components comprised in said combinations can be administered together or separately. The NS3/4a protease inhibitors of formula (I) or any subgroup thereof, and ritonavir or a pharmaceutically acceptable salt or ester thereof, may have dosage levels of the order of 0.02 to 5.0 grams-per-day.

10

When the HCV NS3/4a protease inhibitor of formula (I) and ritonavir are administered in combination, the weight ratio of the HCV NS3/4a protease inhibitor of formula (I) to ritonavir is suitably in the range of from about 40:1 to about 1:15, or from about 30:1 to about 1:15, or from about 15:1 to about 1:15, typically from about 10:1 to about 1:10, and more typically from about 8:1 to about 1:8. Also useful are weight ratios of the HCV NS3/4a protease inhibitors of formula (I) to ritonavir ranging from about 6:1 to about 1:6, or from about 4:1 to about 1:4, or from about 3:1 to about 1:3, or from about 2:1 to about 1:2, or from about 1.5:1 to about 1:1.5. In one aspect, the amount by weight of the HCV NS3/4a protease inhibitors of formula (I) is equal to or greater than that of ritonavir, wherein the weight ratio of the HCV NS3/4a protease inhibitor of formula (I) to ritonavir is suitably in the range of from about 1:1 to about 15:1, typically from about 1:1 to about 10:1, and more typically from about 1:1 to about 8:1. Also useful are weight ratios of the HCV NS3/4a protease inhibitor of formula (I) to ritonavir ranging from about 1:1 to about 6:1, or from about 1:1 to about 5:1, or from about 1:1 to about 4:1, or from about 3:2 to about 3:1, or from about 1:1 to about 2:1 or from about 1:1 to about 1.5:1.

The term "therapeutically effective amount" as used herein means that amount of active compound or component or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought, in the light of the present invention, by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease being treated. Since the instant invention refers to combinations comprising two or more agents, the "therapeutically effective amount" is that amount of the agents taken together so that the combined effect elicits the desired biological or medicinal response. For example, the therapeutically effective amount of a composition comprising (a) the compound of formula (I) and (b) ritonavir, would be the amount of the compound of formula (I) and

35

the amount of ritonavir that when taken together have a combined effect that is therapeutically effective.

In general it is contemplated that an antiviral effective daily amount would be from
5 0.01 mg/kg to 500 mg/kg body weight, more preferably from 0.1 mg/kg to 50 mg/kg body weight. It may be appropriate to administer the required dose as one, two, three, four or more (sub-)doses at appropriate intervals throughout the day. Said (sub-)doses may be formulated as unit dosage forms, for example, containing 1 to 1000 mg, and in particular 5 to 200 mg of active ingredient per unit dosage form.

10

The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight, sex, extent of disorder and general physical condition of the particular patient as well as other medication the individual may be taking, as is
15 well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines.

20

According to one embodiment, the HCV NS3/4a protease inhibitor of formula (I) and ritonavir may be co-administered once or twice a day, preferably orally, wherein the amount of the compounds of formula (I) per dose is from about 1 to about 2500 mg, and the amount of ritonavir per dose is from 1 to about 2500 mg. In another
25 embodiment, the amounts per dose for once or twice daily co-administration are from about 50 to about 1500 mg of the compound of formula (I) and from about 50 to about 1500 mg of ritonavir. In still another embodiment, the amounts per dose for once or twice daily co-administration are from about 100 to about 1000 mg of the compound of formula (I) and from about 100 to about 800 mg of ritonavir. In yet another
30 embodiment, the amounts per dose for once or twice daily co-administration are from about 150 to about 800 mg of the compound of formula (I) and from about 100 to about 600 mg of ritonavir. In yet another embodiment, the amounts per dose for once or twice daily co-administration are from about 200 to about 600 mg of the compound of formula (I) and from about 100 to about 400 mg of ritonavir. In yet another
35 embodiment, the amounts per dose for once or twice daily co-administration are from about 200 to about 600 mg of the compound of formula (I) and from about 20 to about 300 mg of ritonavir. In yet another embodiment, the amounts per dose for once or

twice daily co-administration are from about 100 to about 400 mg of the compound of formula (I) and from about 40 to about 100 mg of ritonavir.

Exemplary combinations of the compound of formula (I) (mg)/ritonavir (mg) for once or twice daily dosage include 50/100, 100/100, 150/100, 200/100, 250/100, 300/100, 350/100, 400/100, 450/100, 50/133, 100/133, 150/133, 200/133, 250/133, 300/133, 50/150, 100/150, 150/150, 200/150, 250/150, 50/200, 100/200, 150/200, 200/200, 250/200, 300/200, 50/300, 80/300, 150/300, 200/300, 250/300, 300/300, 200/600, 400/600, 600/600, 800/600, 1000/600, 200/666, 400/666, 600/666, 800/666, 1000/666, 1200/666, 200/800, 400/800, 600/800, 800/800, 1000/800, 1200/800, 200/1200, 400/1200, 600/1200, 800/1200, 1000/1200, and 1200/1200. Other exemplary combinations of the compound of formula (I) (mg)/ritonavir (mg) for once or twice daily dosage include 1200/400, 800/400, 600/400, 400/200, 600/200, 600/100, 500/100, 400/50, 300/50, and 200/50.

In one embodiment of the present invention there is provided an article of manufacture comprising a composition effective to treat an HCV infection or to inhibit the NS3 protease of HCV; and packaging material comprising a label which indicates that the composition can be used to treat infection by the hepatitis C virus; wherein the composition comprises a compound of the formula (I) or any subgroup thereof, or the combination as described herein.

Another embodiment of the present invention concerns a kit or container comprising a compound of the formula (I) or any subgroup thereof, or a combination according to the invention combining an HCV NS3/4a protease inhibitor of formula (I) or a pharmaceutically acceptable salt thereof, and ritonavir or a pharmaceutically acceptable salt thereof, in an amount effective for use as a standard or reagent in a test or assay for determining the ability of potential pharmaceuticals to inhibit HCV NS3/4a protease, HCV growth, or both. This aspect of the invention may find its use in pharmaceutical research programs.

The compounds and combinations of the present invention can be used in high-throughput target-analyte assays such as those for measuring the efficacy of said combination in HCV treatment.

35

Examples


The following examples are intended to illustrate the present invention and not to limit it thereto.

General: LC/MS analyses were performed on a Waters Alliance 2795 HT attached to a Micromass ZMD mass spectrometer using electrospray ionisation in positive mode.

Eluent: A: water, 0.1% TFA, B: acetonitrile, 0.1% TFA. *Detection*: UV (diode array: 210-300 nm). *Gradients*: Method A: 20 to 70% B in A (1.5 mL min^{-1}) over 5 min.

- 5 Method B: 30 to 80% B in A (1.5 mL min^{-1}) over 5 min. Method C: 40 to 80% B in A (1.5 mL min^{-1}) over 5 min. Method D: 50 to 90% B in A (1.5 mL min^{-1}) over 5 min. Method E: 20 to 70% B in A (0.9 mL min^{-1}) over 2.5 min. Method F: 30 to 80% B in A (0.9 mL min^{-1}) over 2.5 min. Method G: 40 to 80% B in A (0.9 mL min^{-1}) over 2.5 min. Method H: 50 to 90% B in A (0.9 mL min^{-1}) over 2.5 min. *Column*: Methods A-D: 10 Phenomenex, Synergi MAX RP-80A column (5.0 cm, 4.6 mm ϕ , 4 μm). Methods E-H: Phenomenex, Synergi MAX RP-80A column (3.0 cm, 3.0 mm ϕ , 4 μm).

Example 1: Synthesis of 1-[(3-Oxo-2-oxa-bicyclo[2.2.1]heptane-5-carbonyl)-amino]-2-vinyl-cyclopropane carboxylic acid ethyl ester (**3**)




15

To a solution of **1** (857 mg, 5.5 mmol), in DMF (14 mL) and DCM (25 mL) at room temperature, was added **2** (1.15 g, 6.0 mmol), HATU (2.29 g, 6.0 mmol) and DIPEA (3.82 mL, 22 mmol). The reaction was stirred under N_2 -atmosphere at ambient temperature for 1 h. LC/MS analysis showed complete conversion and the reaction mixture was concentrated *in vacuo*. The residue was re-dissolved in DCM (100 mL) and 0.1 M HCl (aq) and the phases separated. The organic phase was washed with NaHCO_3 (aq) and brine, dried (MgSO_4) and filtered. Removal of the solvent *in vacuo* afforded the target compound **3** (1.6 g, 99%). LC/MS (Method A): t_R =2.46 min, >95%, m/z (ESI^+) = 294(MH^+)

25


Example 2: Synthesis of 2-(1-Ethoxycarbonyl-2-vinylcyclopropylcarbamoyl)-4-hydroxy-cyclopentane carboxylic acid diisopropylethylamine salt (**4**)



To a solution of **3** (800 mg, 2.73 mmol) in water (15 mL) in a 20 mL microwave reaction vessel was added DIPEA (1.2 mL, 6.8 mmol) and a stir bar. The reaction vessel was sealed and the immiscible slurry was shaken vigorously before insertion in the microwave oven cavity. After 1 min of pre-stirring, the reaction was irradiated for 40 min to a set temperature of 100 °C. After cooling to 40 °C, the transparent solution was concentrated *in vacuo*, and the residual brown oil co-evaporated 3 times with MeCN to remove any residual water. The crude product **4**, in the form of a DIPEA salt, was immediately taken forward to the next step. LC/MS (Method A): t_R =1.29 min, >95%, m/z (ESI⁺)= 312(MH⁺).


10

Example 3: Synthesis of 1-{[2-(Hex-5-enylmethylcarbamoyl)-4-hydroxycyclopentane carbonyl]amino}-2-vinylcyclopropane carboxylic acid ethyl ester (**6**)




The crude compound **4** (5.5 mmol) was dissolved in DCM (50 mL) and DMF (14 mL) followed by addition of HATU (2.09 g, 5.5 mmol), **5** (678 mg, 6.0 mmol) and DIPEA (3.08 mL, 17.5 mmol) at room temperature. The reaction was stirred at ambient temperature for 1 h. LC/MS analysis showed complete conversion and the reaction mixture was concentrated *in vacuo*. The residue was re-dissolved in EtOAc (100 mL) and the organic phase washed with 0.1 M HCl (aq), K₂CO₃ (aq) and brine, dried (MgSO₄) and filtered. Evaporation of the solvent *in vacuo* gave an oil which was purified by flash chromatography (Silica, EtOAc:MeOH) to afford the target compound **6** (1.65 g, 74%). TLC (Silica): MeOH:EtOAc 5:95, R_f =0.5; LC/MS (Method A): t_R =3.44 min, >95%, m/z (ESI⁺)= 407(MH⁺).

Example 4: Synthesis of 1-{[2-(Hex-5-enylmethylcarbamoyl)-4-hydroxycyclopentane-carbonyl]amino}-2-vinylcyclopropanecarboxylic acid (**7**).



Compound **6** (493 mg, 1.21 mmol) was dissolved in DMF (1 mL) and transferred to a 20 mL microwave reaction vessel. Then, aqueous LiOH (2 M, 10.5 mL) and a stirbar were added. The reaction vessel was sealed and the immiscible slurry was shaken
 5 vigorously before insertion in the microwave cavity. The reaction was irradiated for 30 min to 130 °C. The reaction mixture was cooled to 40 °C and the clear solution acidified to pH 2 with aqueous HCl (1 M, 24 mL) and extracted 3 times with EtOAc (20 mL). The pooled organic layers were washed with brine, dried (MgSO₄) and filtered. The solvent was evaporated *in vacuo* to afford compound **7** (410 mg, 90 %).
 10 LC/MS (Method A): t_R = 2.46 min, >95%, m/z (ESI⁺) = 379(MH⁺).


Example 5: Synthesis of 4-Hydroxy-cyclopentane-1,2-dicarboxylic acid 1-[(1-cyclopropanesulfonylamino carbonyl-2-vinyl-cyclopropyl)-amide] 2-(hex-5-enyl-methyl-amide) (**8**).




15 The crude acid **7** (410 mg, 1.09 mmol) was dissolved in DMF (1.5 mL) and DCM (4.5 mL) followed by addition of EDAC (417 mg, 2.18 mmol) at room temperature. The mixture was allowed to incubate with stirring at room temperature. After 10 min, DMAP (133 mg, 1.09 mmol) was added followed by another 20 min incubation at
 20 room temperature. Subsequently, a pre-mixed solution of cyclopropanesulfonic acid amide (527 mg, 4.36 mmol) and DBU (663 mg, 4.36 mmol) in DMF (2 mL) and DCM (2 mL) was added followed by heating in the microwave oven to 100°C for 30 min. The resulting red solution was concentrated *in vacuo* and re-dissolved in EtOAc (20 mL). The organic phase was washed with 1 M HCl (aq) (3x 10 mL) and brine
 25 (10 mL), dried (MgSO₄) and filtered. The solvent was evaporated *in vacuo* to yield the

crude sulfonamide which was further purified by chromatography (Silica, EtOAc:MeOH, 97.5:2.5) to afford the target compound **8** (403 mg, 77%); LC/MS (Method A): t_R = 3.31 min, >95%, m/z (ESI^+) = 482(MH^+).

- 5 **Example 6-1:** Synthesis of 2,3-Dihydroindole-1-carboxylic acid 3-(1-cyclopropane-sulfonylaminocarbonyl-2-vinylcyclopropylcarbamoyl)-4-(hex-5-enylmethylcarbamoyl)cyclopentyl ester (**11**).




- 10 Compound **8** (19.4 mg, 40 μ mol) was dissolved in DCM (1.8 mL) followed by addition of solid $NaHCO_3$ (14 mg, 160 μ mol) and a stirbar. To this slurry was then added phosgene in toluene (1.93 M, 430 μ l, 0.8 mmol) and the mixture was stirred vigorously for 2 h to afford the chloroformate **9**. LC/MS (Method G): t_R = 2.65 min, >95%, m/z (ESI^+) = 544(MH^+). The solvent was evaporated *in vacuo* and the residue was co-
- 15 evaporated 3 times with DCM to remove any residual phosgene.
- The afforded chloroformate **9** was subsequently re-dissolved in DCM (1 mL) and 2,3 dihydroindole (68 μ mol) was added. The mixture was allowed to stir at ambient temperature for 2 h after which time LC/MS showed complete conversion. Then, DCM (1 mL) was added to the mixture and the solution was washed 2 times with 1 M HCl
- 20 (aq), $NaHCO_3$ (aq) and brine. The organic phase was dried ($MgSO_4$) and filtered. Evaporation of the solvent *in vacuo* gave a crude which was further purified by preparative LC/MS to afford compound **10**: LC/MS (Method H): t_R = 1.58 min, >95%, m/z (ESI^+) = 627(MH^+).
- 25 **Example 6-2:** 3,4-Dihydro-2*H*-quinoline-1-carboxylic acid 3-(1-cyclopropanesulfonylaminocarbonyl-2-vinylcyclopropylcarbamoyl)-4-(hex-5-enylmethylcarbamoyl)cyclopentyl ester (**11**).



The title compound was synthesized from 1,2,3,4-tetrahydroquinoline according to the procedure described in Example 6-1. LC/MS (Method H): t_R = 1.74 min, >95%, m/z (ESI^+) = 641(MH^+).


5

Example 6-3: 2,3-Dihydrobenzo[1,4]oxazine-4-carboxylic acid 3-(1-cyclopropane sulfonylaminocarbonyl-2-vinylcyclopropylcarbamoyl)-4-(hex-5-enylmethylcarbamoyl) cyclopentyl ester (**12**).



10 The title compound was synthesized from 3,4-dihydro-2H-benzo[1,4]oxazine according to the procedure described in Example 6-1. LC/MS (Method H): t_R = 1.56 min, >95%, m/z (ESI^+) = 643(MH^+).


15 **Example 6-4:** 1,3-Dihydroisindole-2-carboxylic acid 3-(1-cyclopropanesulfonyl aminocarbonyl-2-vinylcyclopropylcarbamoyl)-4-(hex-5-enylmethylcarbamoyl) cyclopentyl ester (**13**)



The title compound was synthesized from 2,3-dihydro-1H-isoindole according to the procedure described in Example 6-1. LC/MS (Method H): t_R = 1.37 min, >95%, m/z (ESI^+) = 627(MH^+).

5


Example 7: 3,4-Dihydro-1H-isoquinoline-2-carboxylic acid 3-(1-cyclopropane sulfonylaminocarbonyl-2-vinylcyclopropylcarbamoyl)-4-(hex-5-enylmethyl carbamoyl)cyclopentyl ester (**16**).



- 10 *p*-Nitrophenyl chloroformate (25.9 mg, 0.129 mmol) was dissolved in MeCN (1 mL). To this solution was added solid $NaHCO_3$ (15.7 mg, 0.19 mmol) and the suspension was cooled in an ice/water bath. To the cooled solution was then added a solution of 1,2,3,4-tetrahydro-isoquinoline (0.123 mmol) in MeCN (0.5 mL) and the reaction was allowed to incubate at ambient temperature for 2 h. LC/MS analysis showed complete
- 15 conversion to compound 15. This solution was then added to a mixture of 8 (49.2 mg, 102 μ mol) and NaH (60 % in oil) (4.5 mg, 112 μ mol) followed by heating of the reaction to 50 °C for 1 h. The reaction was quenched with NH_4Cl (aq) (5 mL) and EtOAc (5 mL) was added. The organic layer was washed with 1 M HCl (aq) and brine, dried ($MgSO_4$) and filtered. Evaporation of the solvent gave an oil which was further


purified using preparative LC/MS to afford the target product **16**: LC/MS (Method X): $t_R = 5.13$ min, >90%, m/z (ESI^+) = 641 (MH^+).

Example 8-1: 2,3-Dihydroindole-1-carboxylic acid 4-cyclopropanesulfonylamino carbonyl-13-methyl-2,14-dioxo-3,13-diazatricyclo[13.3.0.0^{4,6}]octadec-7-en-17-yl ester, (**17**).



Compound **10** (14.6 μ mol) was dissolved in DCE (dried over mol sieves, N_2 -gassed) (10 mL) in a 20 mL microwave reaction vessel with a stir bar. To this solution was added Hoveyda-Grubb's 2nd generation catalyst (2.3 mg, 3.6 μ mol) and the reaction vessel was purged with $N_2(g)$ and sealed. The reaction was irradiated for 15 min with a set temperature of 150 $^{\circ}C$. The solvent was removed *in vacuo* and the residue purified by flash chromatography (Silica; DCM, then 10% MeOH in DCM). The product was subsequently purified by preparative LC/MS to afford the target compound **17**: LC/MS (Method H): $t_R = 1.13$ min, >95%, m/z (ESI^+) = 599(MH^+).

Example 8-2: 3,4-Dihydro-1*H*-isoquinoline-2-carboxylic acid 4-cyclopropanesulfonyl amino carbonyl-13-methyl-2,14-dioxo-3,13-diaza-tricyclo[13.3.0.0^{4,6}]octadec-7-en-17-yl ester (**18**)




20

The title compound was prepared from 3,4-Dihydro-1*H*-isoquinoline-2-carboxylic acid 3-(1-cyclopropane sulfonylamino carbonyl-2-vinylcyclopropyl carbamoyl)-4-(hex-5-


enylmethyl carbamoyl)cyclopentyl ester (**16**) following the procedure described in Example 8-1. LC/MS (Method A): $t_R = 4.51$ min, >95%, m/z (ESI^+) = 613(MH^+).

Example 8-3: 2,3-Dihydrobenzo[1,4]oxazine-4-carboxylic acid 4-cyclopropane sulfonylamino carbonyl-13-methyl-2,14-dioxo-3,13-diazatricyclo[13.3.0.0^{4,6}]octadec-7-en-17-yl ester (**19**)




The title compound was prepared from 2,3-Dihydrobenzo[1,4]oxazine-4-carboxylic acid 3-(1-cyclopropane sulfonylamino carbonyl-2-vinylcyclopropylcarbamoyl)-4-(hex-5-enylmethyl carbamoyl) cyclopentyl ester (**12**) following the procedure described in Example 8-1: LC/MS (Method H): $t_R = 1.11$ min, >95%, m/z (ESI^+) = 615(MH^+).

Example 8-4: 1,3-Dihydro-isoindole-2-carboxylic acid 4-cyclopropanesulfonylamino carbonyl-13-methyl-2,14-dioxo-3,13-diaza-tricyclo[13.3.0.0^{4,6}]octadec-7-en-17-yl ester, (**20**).




The title compound was prepared from 1,3-Dihydroisoindole-2-carboxylic acid 3-(1-cyclopropanesulfonyl aminocarbonyl-2-vinylcyclopropylcarbamoyl)-4-(hex-5-enylmethyl carbamoyl) cyclopentyl ester (**13**) following the procedure described in Example 8-1: LC/MS (Method F): $t_R = 2.33$ min, >95%, m/z (ESI^+) = 599(MH^+).

Example 8-5: 3,4-Dihydro-2*H*-quinoline-1-carboxylic acid 4-cyclopropane sulfonyl-aminocarbonyl-13-methyl-2,14-dioxo-3,13-diaza-tricyclo[13.3.0.0^{4,6}]octadec-7-en-17-yl ester, (**21**):




- 5 The title compound was prepared from 3,4-Dihydro-2*H*-quinoline-1-carboxylic acid 3-(1-cyclopropanesulfonyl aminocarbonyl-2-vinylcyclopropylcarbamoyl)-4-(hex-5-enylmethyl carbamoyl) cyclopentyl ester (**11**) following the procedure described in Example 8-1: LC/MS (Method H): t_R = 1.25 min, >95%, m/z (ESI^+) = 613(MH^+).

10 **Example 9**




2-(1-Ethoxycarbonyl-2-vinyl-cyclopropylcarbamoyl)-4-hydroxy-pyrrolidine-1-carboxylic acid tert-butyl ester (22)

- Boc-protected 4-hydroxy proline (4 g, 17.3 mmol), HATU (6.9 g, 18.2 mmol) and 1-amino-2-vinyl-cyclopropanecarboxylic acid ethyl ester prepared as described in WO03/099274, (3.5 g, 18.3 mmol) were dissolved in DMF (60 ml) and cooled to 0 ° on an ice-bath. Diisopropylethyl amine (DIPEA) (6ml) was added. The ice-bath was removed and the mixture was left at ambient temperature over-night. Dichloromethane (~80 ml) was then added and the organic phase was washed with aqueous sodium hydrogen carbonate, citric acid, water, brine and dried over sodium sulphate.
- 20 Purification by flash chromatography (ether → 7% methanol in ether) gave pure title compound (6.13 g, 96%)

Example 10


2-(1-Ethoxycarbonyl-2-vinyl-cyclopropylcarbamoyl)-4-(4-nitro-benzoyloxy)-pyrrolidine-1-carboxylic acid tert-butyl ester (23)

- 5 Compound 22 (from example 9) (11.8 g, 32.0 mmol) and pyridine (27 ml, 305 mmol) was dissolved in DCM (200 ml) and cooled to 0°C, 4-nitrobenzoyl chloride (6.6 g, 35.6 mmol) was added and the solution was stirred at room temperature overnight. The reaction mixture was washed with NaHCO₃ (aq), aqueous citric acid and brine, dried over MgSO₄ and evaporated on silica. The crude product was purified by column
- 10 chromatography on silica (EtOAc/n-Heptane: 50/50) to give 11.84 g, 72 % of the title compound **5**.

Example 11


- 15 4-Nitro-benzoic acid 5-(1-ethoxycarbonyl-2-vinyl-cyclopropylcarbamoyl)-pyrrolidin-3-yl ester (24)

Compound 23 (11.84 g, 22.9 mmol) was deprotected in TFA (30 ml) dissolved in DCM (100 ml) and then worked up by methods known in the chemical art to give the title compound (9.37 g, 98 %).

Example 12

4-Nitro-benzoic acid 5-(1-ethoxycarbonyl-2-vinyl-cyclopropylcarbamoyl)-1-[hept-6-enyl-(4-methoxy-benzyl)-carbamoyl]-pyrrolidin-3-yl ester (25)


- 5 Compound 24 (4.68 g, 11.2 mmol) was dissolved in THF (100 ml), NaHCO₃ (s) (appr. 5 ml) was added followed by phosgene-solution (20 % in toluene, 11.6 ml, 22.5 mmol). The reaction mixture was stirred vigorously for 1h and then filtrated, evaporated and redissolved in DCM (100ml). NaHCO₃ (s) (appr. 5 ml) was added followed by hept-6-enyl-(4-methoxy-benzyl)-amin (3.92 g, 16.8 mmol). The reaction mixture was stirred at
- 10 room temperature overnight, filtrated and evaporated on silica. The crude product was purified by column chromatography on silica (EtOAc/ n-Heptane: 25/75) to give the title compound (6.9 g, 91 %).

Example 13

15


14-(4-Methoxy-benzyl)-18-(4-nitro-benzoyloxy)-2,15-dioxo-3,14,16-triazatricyclo[14.3.0.0*4,6*]nonadec-7-ene-4-carboxylic acid ethyl ester (26)

- Compound 25 (406 mg, 0.6 mmol) was dissolved in DCE (250 ml) and degassed. Hoveyda-Grubbs Catalyst 2nd generation (26 mg, 0.042 mmol) was added and the
- 20 solution was heated to reflux. After 3 h the solution was evaporated and used direct in the next step.

Example 14


18-Hydroxy-14-(4-methoxy-benzyl)-2,15-dioxo-3,14,16-triaza-tricyclo[14.3.0.0*4,6*]-nonadec-7-ene-4-carboxylic acid ethyl ester (27)

- 5 Crude compound 26 (445 mg) was dissolved in THF (20 ml), MeOH (10 ml) and water (10 ml). After cooling to 0°C 1M LiOH (2 ml) was added. After 1.5 h the hydrolysis was completed and HOAc (1ml) was added and the solution was evaporated to approx 10 ml. Water was added and the mixture was extracted with DCM (2 x 30 ml). The pooled organic phase was washed with NaHCO₃ (aq), water, brine and dried over
- 10 MgSO₄. The crude product was purified by column chromatography on silica (DCM/MeOH: 100/0 – 80/20) to give the title compound (201 mg, 67 %).

Example 15

- 15 18-Ethoxymethoxy-14-(4-methoxy-benzyl)-2,15-dioxo-3,14,16-triaza-tricyclo[14.3.0.0*4,6*]-nonadec-7-ene-4-carboxylic acid ethyl ester (28)


- To a stirred solution of the alcohol 27 (1.35 g, 2.70 mmol, 75 % purity) and N-ethyl-diisopropylamine (1.42 ml, 8.1 mmol) in dichloromethane (15 ml) at 0 °C was added chloromethyl ethyl ether (0.5 ml, 5.4 mmol). After stirring at rt on the reaction mixture
- 20 was cooled to 0 °C and more N-ethyldiisopropylamine (1 ml, 5.7 mmol) and chloromethyl ethyl ether (0.3 ml, 3.2 mmol) was added, then stirred additional 16 h at rt. The reaction mixture was then directly applied on a silicagel column and eluted using stepwise gradient elution (ethyl acetate in hexane 50-80 %). Concentration of the appropriate fractions gave the title compound as a slight brown syrup which
- 25 crystallized upon standing (0.8 g, 53%). LR-MS: Calcd for C₃₀H₄₄N₃O₇: 558. Found: 558 [M+H].

Example 16


18-Ethoxymethoxy-14-(4-methoxy-benzyl)-2,15-dioxo-3,14,16-triazatricyclo[14.3.0.0*4,6*]nonadec-7-ene-4-carboxylic acid (29)

- 5 A solution of the ester 28 (0.775 g, 1.39 mmol) in 1:1:1 THF-Methanol-aq. 1M LiOH (36 ml) was stirred at rt for 3.5 h after which TLC (95:5 and 9:1 dichloromethane-methanol) and LC-MS indicated complete conversion into the carboxylic acid. The reaction mixture was then concentrated into approximately 1/3 of the volume, then diluted with water (10 ml) and acidified to approx. pH 4 using aq. 10 % citric acid
- 10 (60 ml) upon which a precipitate formed. The mixture was washed with ethyl acetate (3 x 25 ml) and the combined organic layers were washed with brine (2 x 50 ml), then dried (Na₂SO₄), filtered and concentrated. The residue was concentrated from toluene (3 x 10 ml) which gave the crude title compound as an off-white foam (0.75 g, quantitative). LR-MS: Calcd for C₂₈H₄₀N₃O₇: 530. Found: 530 [M-H].

15

Example 17Compound 30


- To a solution of the carboxylic acid 29 (approx. 1.39 mmol) in dichloromethane (10 ml) at rt was added N-Ethyl-N'-(3-dimethylaminopropyl)carbodiimide x HCl (0.32 g, 1.67 mmol), then stirred overnight after which LC-MS indicated complete conversion of the acid into the product. The reaction mixture was then diluted with dichloromethane (10 ml), washed with water (3 x 10 ml), then dried (Na₂SO₄) filtered and concentrated into a colorless solid (crude yield: 0.7 g) which was used immediately
- 25 in the next step. LR-MS: Calcd for C₂₈H₃₈N₃O₆: 512. Found: 512 [M+H].

Example 18

Cyclopropanesulfonic acid [18-ethoxymethoxy-14-(4-methoxy-benzyl)-2,15-dioxo-3,14,16-triaza-tricyclo[14.3.0.0*4,6*]nonadec-7-ene-4-carbonyl]-amide (31)

- 5 To a stirred solution of the crude oxazolinone 30 (0.328 g, 0.64 mmol) in dichloromethane (4 ml) was added cyclopropylsulfonamide (0.117 g, 0.96 mmol) and 1,8-diazabicyclo[5.4.0]-undec-7-ene (0.19 ml, 1.3 mmol), then stirred at rt overnight. The reaction mixture was monitored by LC-MS then diluted with dichloromethane (20 ml), washed successively with aq. 10 % citric acid (3 x 15 ml) and brine (1 x 15 ml), then dried (Na₂SO₄), filtered and concentrated into an off-white foam. Column chromatography of the residue using stepwise gradient elution (ethyl acetate in toluene 60-100 %) followed by concentration and drying of the appropriate fractions gave the title compound as a colorless foam (0.27 g, 66 % over 3 steps).
- 15 NMR data (500 MHz, DMSO-d₆): ¹H, 0.9-1.6 (m, 14H), 1.80 (m, 1H), 1.90 (m, 1H), 2.0-2.2 (m, 3H), 2.25 (m, 1H), 2.95 (m, 1H), 3.05 (m, 1H), 3.3-3.4 (m, 2H), 3.50 (q, 2H), 3.7-3.8 (m, 4H), 3.97 (d, 1H), 4.3-4.4 (m, 2H), 4.55 (d, 1H), 4.63 (m, 2H), 5.12 (m, 1H), 5.70 (m, 1H), 6.88 (d, 2H), 7.19 (d, 2H), 8.12 (s, 1H). LR-MS: Calcd for C₃₁H₄₅N₄O₈S: 633. Found: 633 [M+H].

20 Example 19




Cyclopropanesulfonic acid (18-hydroxy-2,15-dioxo-3,14,16-triaza-tricyclo[14.3.0.0*4,6*]nonadec-7-ene-4-carbonyl)-amide (32)

- 25 A solution of the acetal 31 (0.038 g, 0.06 mmol) in 1:1:1 THF-methanol-2 M aq. hydrochloric acid (1.5 ml) was stirred at rt for 30 min, then additional conc. hydrochloric acid (0.1 ml) was added and then stirred at rt overnight. The reaction mixture was then neutralized using aq. saturated sodium hydrogen carbonate, then concentrated onto silica. Flash chromatography of the residue using 9:1 ethyl acetate-

methanol gave a colorless foam (0.020 g, 73 %). LR-MS: Calcd for C₂₀H₂₉N₄O₆S: 453. Found: 453 [M-H].

Example 20-1




5

1,3-Dihydro-isoindole-2-carboxylic acid 4-cyclopropanesulfonylaminocarbonyl-2,15-dioxo-3,14,16-triaza-tricyclo[14.3.0.0*4,6*]nonadec-7-en-18-yl ester (33)

Alcohol 32 (25 mg, 55 μ mol) was dissolved in dry DCM (2 mL). To this solution was added solid NaHCO₃ (14 mg, 165 μ mol) and phosgene (1.9 M in toluene, 868 μ L, 1.65 mmol). The mixture was stirred for 48 h to afford the intermediate chloroformate. LC/MS (Method F): t_R = 2.32 min, m/z (ESI⁺) = 516 (MH⁺). The solvent was removed *in vacuo* and the residue was co-evaporated with DCM to remove any residual phosgene. The afforded chloroformate was subsequently re-dissolved in dry DCE (2 ml) and isoindoline (83 μ mol) was added followed by solid K₂CO₃ (110 μ mol) and powdered 4Å mol.sieves (1 spatula). The mixture was heated to 100 °C for 45 min, after which time LC/MS analysis showed no remaining chloroformate. The reaction was filtered and the filtrate concentrated *in vacuo* to afford a crude which was purified by preparative LC/MS to yield the title compound. LC/MS (Method H): t_R = 1.55 min, >95%, m/z (ESI⁺) = 600 (MH⁺).

20


Example 20-2



2,3-Dihydro-indole-1-carboxylic acid 4-cyclopropanesulfonylaminocarbonyl-2,15-dioxo-3,14,16-triaza-tricyclo[14.3.0.0*4,6*]nonadec-7-en-18-yl ester (34)

The title compound was prepared according to the procedure described in Example 20-1, except that indoline was used instead of isoindoline. LC/MS (Method H): t_R =
 5 1.68 min, 95%, m/z (ESI^+)= 600 (MH^+).

Example 20-3




3,4-Dihydro-1H-isoquinoline-2-carboxylic acid 4-cyclopropanesulfonylaminocarbonyl-2,15-dioxo-3,14,16-triaza-tricyclo[14.3.0.0*4,6*]nonadec-7-en-18-yl ester (35)

The title compound was prepared according to the procedure described in Example 20-1, except that 1,2,3,4-tetrahydro-isoquinoline was used instead of isoindoline. LC/MS (Method H): t_R = 1.60 min, 95%, m/z (ESI^+)= 614 (MH^+).

15


Example 20-4



3,4-Dihydro-2H-quinoline-1-carboxylic acid 4-cyclopropanesulfonylaminocarbonyl-2,15-dioxo-3,14,16-triaza-tricyclo[14.3.0.0*4,6*]nonadec-7-en-18-yl ester (36)


The title compound was prepared according to the procedure described in Example 20-1, except that 1,2,3,4-tetrahydro-quinoline was used instead of isoindoline. LC/MS (Method H): t_R = 1.77 min, 95%, m/z (ESI^+)= 614 (MH^+).

20

Example 20-5

5-Methyl-2,3-dihydro-indole-1-carboxylic acid 4-cyclopropanesulfonylamino-carbonyl-2,15-dioxo-3,14,16-triaza-tricyclo[14.3.0.0*4,6*]nonadec-7-en-18-yl ester (37)

The title compound was prepared according to the procedure described in Example 20-1, except that 5-methyl-2,3-dihydro-1H-indole was used instead of isoindoline. LC/MS (Method H): t_R = 1.91 min, 95%, m/z (ESI^+) = 614 (MH^+).


10 Example 20-6

5-Dimethylsulfonyl-2,3-dihydro-indole-1-carboxylic acid 4-cyclopropanesulfonylaminocarbonyl-2,15-dioxo-3,14,16-triaza-tricyclo[14.3.0.0*4,6*]nonadec-7-en-18-yl ester (38)

15 The title compound was prepared according to the procedure described in Example 20-1, except that 2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide was used instead of isoindoline. LC/MS (Method H): t_R = 1.53 min, 95%, m/z (ESI^+) = 707 (MH^+).

20 Example 21: Synthesis of crystalline cyclopentane

Synthesis of 3-Oxo-2-oxa-bicyclo[2.2.1]heptane-5-carboxylic acid *tert*-butyl ester (40)




DMAP (14 mg, 0.115 mmol) and Boc_2O (252 mg, 1.44 mmol) was added to a stirred solution of **39** (180 mg, 1.15 mmol) in 2 mL CH_2Cl_2 under inert argon atmosphere at 0°C . The reaction was allowed to warm to room temperature and was stirred overnight.

5 The reaction mixture was concentrated and the crude product was purified by flash column chromatography (toluene/ethyl acetate gradient 15:1, 9:1, 6:1, 4:1, 2:1) which gave the title compound (124 mg, 51%) as white crystals.

$^1\text{H-NMR}$ (300 MHz, CD_3OD) δ 1.45 (s, 9H), 1.90 (d, $J = 11.0$ Hz, 1H), 2.10-2.19 (m, 3H), 2.76-2.83 (m, 1H), 3.10 (s, 1H), 4.99 (s, 1H); $^{13}\text{C-NMR}$ (75.5 MHz, CD_3OD) δ 27.1, 33.0, 37.7, 40.8, 46.1, 81.1, 81.6, 172.0, 177.7.

Alternative method for the preparation of compound **40**



15 Compound **39** (13.9 g, 89 mmol) was dissolved in dichloromethane (200 ml) and then cooled to approximately -10°C under nitrogen. Isobutylene was then bubbled into the solution until the total volume had increased to approximately 250 mL which gave a turbid solution. $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (5.6 ml, 44.5 mmol, 0.5 eq.) was added and the reaction mixture was kept at approximately -10°C under nitrogen. After 10 min, a clear solution

20 was obtained. The reaction was monitored by TLC (EtOAc-Toluene 3:2 acidified with a few drops of acetic acid and hexane-EtOAc 4:1, staining with basic permanganate solution). At 70 min only traces of compound **39** remained and aq. saturated NaHCO_3 (200 ml) was added to the reaction mixture, which was then stirred vigorously for 10 min. The organic layer was washed with saturated NaHCO_3 (3 x 200 ml) and brine

25 (1 x 150 ml), then dried with sodium sulfite, filtered and the residue was evaporated to an oily residue. Upon addition of hexane to the residue, the product precipitated. Addition of more hexane and heating to reflux gave a clear solution from which the product crystallized. The crystals were collected by filtration and were washed with hexane (rt), then air-dried for 72 h giving colourless needles (12.45 g, 58.7 mmol,

30 66%).

Example 22: Activity of compounds of formula (I)

Replicon assay

The compounds of formula (I) were examined for activity in the inhibition of HCV RNA replication in a cellular assay. The assay demonstrated that the compounds of formula (I) exhibited activity against HCV replicons functional in a cell culture. The cellular assay was based on a bicistronic expression construct, as described by Lohmann et al. (1999) Science vol. 285 pp. 110-113 with modifications described by Krieger et al. (2001) Journal of Virology 75: 4614-4624, in a multi-target screening strategy. In essence, the method was as follows.

- 5 The assay utilized the stably transfected cell line Huh-7 luc/neo (hereafter referred to as Huh-Luc). This cell line harbors an RNA encoding a bicistronic expression construct comprising the wild type NS3-NS5B regions of HCV type 1b translated from an Internal Ribosome Entry Site (IRES) from encephalomyocarditis virus (EMCV), preceded by a reporter portion (FfL-luciferase), and a selectable marker portion (neo^R, neomycine phosphotransferase). The construct is bordered by 5' and 3' NTRs (non-translated regions) from HCV type 1b. Continued culture of the replicon cells in the presence of G418 (neo^R) is dependent on the replication of the HCV RNA. The stably transfected replicon cells that express HCV RNA, which replicates autonomously and to high levels, encoding *inter alia* luciferase, are used for screening the antiviral compounds.
- 10
15
20

The replicon cells were plated in 384 well plates in the presence of the test and control compounds which were added in various concentrations. Following an incubation of three days, HCV replication was measured by assaying luciferase activity (using standard luciferase assay substrates and reagents and a Perkin Elmer ViewLux[™] ultraHTS microplate imager). Replicon cells in the control cultures have high luciferase expression in the absence of any inhibitor. The inhibitory activity of the compound on luciferase activity was monitored on the Huh-Luc cells, enabling a dose-response curve for each test compound. EC50 values were then calculated, which value represents the amount of the compound required to decrease by 50% the level of detected luciferase activity, or more specifically, the ability of the genetically linked HCV replicon RNA to replicate.

25
30

Inhibition assay


- 35 The aim of this *in vitro* assay was to measure the inhibition of HCV NS3/4A protease complexes by the compounds of the present invention. This assay provides an indication of how effective compounds of the present invention would be in inhibiting HCV NS3/4A proteolytic activity.

The inhibition of full-length hepatitis C NS3 protease enzyme was measured essentially as described in Poliakov, 2002 Prot Expression & Purification 25 363 371. Briefly, the hydrolysis of a depsipeptide substrate, Ac-DED(Edans)EEAbuψ[COO]ASK(Dabcyl)-NH₂ (AnaSpec, San José, USA), was measured spectrofluorometrically in the presence of a peptide cofactor, KKGSVVIVGRIVLSGK (Åke Engström, Department of Medical Biochemistry and Microbiology, Uppsala University, Sweden). [Landro, 1997 #Biochem 36 9340-9348]. The enzyme (1 nM) was incubated in 50 mM HEPES, pH 7.5, 10 mM DTT, 40% glycerol, 0.1% n-octyl-D-glucoside, with 25 μM NS4A cofactor and inhibitor at 30 °C for 10 min, whereupon the reaction was initiated by addition of 0.5 μM substrate. Inhibitors were dissolved in DMSO, sonicated for 30 sec. and vortexed. The solutions were stored at - 20°C between measurements.






The final concentration of DMSO in the assay sample was adjusted to 3.3%. The rate of hydrolysis was corrected for inner filter effects according to published procedures. [Liu, 1999 Analytical Biochemistry 267 331-335]. K_i values were estimated by non-linear regression analysis (GraFit, Erithacus Software, Staines, MX, UK), using a model for competitive inhibition and a fixed value for K_m (0.15 μM). A minimum of two replicates was performed for all measurements.

20

The following Table 1 lists compounds that were prepared according to any one of the above examples. The activities of the compounds tested are also depicted in Table 1.




Compound nr.		n	EC ₅₀ (μM) Replicon assay	K _i (nM) Enzymatic assay
1		4	0.809	5.1


Compound nr.		n	EC ₅₀ (μM) Replicon assay	Ki (nM) Enzymatic assay
2		4	3.369	14.7
3		4	>10	62
5		4	>10	12
6		4	>10	24

Claims

1. A compound having the formula



- 5 an *N*-oxide, salt, or stereoisomer thereof,
 wherein each dashed line (represented by - - - -) represents an optional double bond;
X is N, CH and where **X** bears a double bond it is C;
R¹ is -OR⁶, -NH-SO₂R⁷;
R² is hydrogen, and where **X** is C or CH, **R**² may also be C₁₋₆alkyl;
 10 **R**³ is hydrogen, C₁₋₆alkyl, C₁₋₆alkoxyC₁₋₆alkyl, or C₃₋₇cycloalkyl;
n is 3, 4, 5, or 6;
R⁴ and **R**⁵ taken together with the nitrogen atom to which they are attached form a bicyclic ring system selected from




- 15 wherein said ring system may optionally be substituted with one, two or three substituents independently selected from halo, hydroxy, oxo, nitro, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₁₋₆alkoxycarbonyl, amino, azido, mercapto, polyhaloC₁₋₆alkyl;
R⁶ is hydrogen; aryl; Het; C₃₋₇cycloalkyl optionally substituted with C₁₋₆alkyl; or
 20 C₁₋₆alkyl optionally substituted with C₃₋₇cycloalkyl, aryl or with Het;

R⁷ is aryl; Het; C₃₋₇cycloalkyl optionally substituted with C₁₋₆alkyl; or C₁₋₆alkyl optionally substituted with C₃₋₇cycloalkyl, aryl or with Het;


aryl as a group or part of a group is phenyl or naphthyl, each of which may be optionally substituted with one, two or three substituents selected from halo, hydroxy, nitro, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkyl, C₁₋₆alkylcarbonyl, amino, mono- or diC₁₋₆alkylamino, azido, mercapto, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkoxy, C₃₋₇cycloalkyl, pyrrolidinyl, piperidinyl, piperazinyl, 4-C₁₋₆alkyl-piperazinyl, 4-C₁₋₆alkylcarbonyl-piperazinyl, and morpholinyl; wherein the morpholinyl and piperidinyl groups may be optionally substituted with one or with two C₁₋₆alkyl radicals;

Het as a group or part of a group is a 5 or 6 membered saturated, partially unsaturated or completely unsaturated heterocyclic ring containing 1 to 4 heteroatoms each independently selected from nitrogen, oxygen and sulfur, and being optionally substituted with one, two or three substituents each independently selected from the group consisting of halo, hydroxy, nitro, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkyl, C₁₋₆alkylcarbonyl, amino, mono- or di-C₁₋₆alkylamino, azido, mercapto, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkoxy, C₃₋₇cycloalkyl, pyrrolidinyl, piperidinyl, piperazinyl, 4-C₁₋₆alkyl-piperazinyl, 4-C₁₋₆alkylcarbonyl-piperazinyl, and morpholinyl; wherein the morpholinyl and piperidinyl groups may be optionally substituted with one or with two C₁₋₆alkyl radicals.


2. A compound according to claim 1, wherein the compound has the formula (I-c), (I-d), or (I-e):




(I-c)



(I-d)




3. A compound according to any one of claims 1-2, wherein R^4 and R^5 taken together with the nitrogen atom to which they are attached form a bicyclic ring system selected from




- wherein the phenyl of said bicyclic ring system is optionally substituted with one or two substituents independently selected from halo, hydroxy, cyano, carboxyl, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkoxy-carbonyl, amino, and polyhalo C_{1-6} alkyl.

4. A compound according to any one of claims 1-2, wherein R^4 and R^5 taken together with the nitrogen atom to which they are attached form a bicyclic ring system selected from




wherein the pyrrolidine, piperidine, or morpholine rings of said bicyclic ring system are optionally substituted with one or two substituents independently selected from C₁₋₆alkyl, C₁₋₆alkoxy, and C₁₋₆alkoxyC₁₋₆alkyl.


- 5 5. A compound according to any one of claims 1-4, wherein
 (a) R¹ is -OR⁶, wherein R⁶ is C₁₋₆alkyl or hydrogen; or
 (b) R¹ is -NHS(=O)₂R⁷, wherein R⁷ is methyl, cyclopropyl, or phenyl.
6. A compound according to any of claims 1-5 other than an N-oxide, or salt.
- 10 7. A combination comprising
 (a) a compound as defined in any one of claims 1 to 6 or a pharmaceutically acceptable salt thereof; and
 (b) ritonavir, or a pharmaceutically acceptable salt thereof.
- 15 8. A pharmaceutical composition comprising a carrier, and as active ingredient an anti-virally effective amount of a compound as claimed in any one of claims 1-6 or a combination according to claim 7.
- 20 9. A compound according to any of claims 1-6 or a combination according to claim 7, for use as a medicament.
10. Use of a compound according to any of claims 1-6 or a combination according to claim 7, for the manufacture of a medicament for inhibiting HCV replication.
- 25 11. A method of inhibiting HCV replication in a warm-blooded animal said method comprising the administration of an effective amount of a compound according to any of claims 1-6 or an effective amount of each component of the combination according to claim 7.
- 30 12. A process for preparing a compound as claimed in any of claims 1 - 6, wherein said process comprises:
- 35 (a) preparing a compound of formula (I) wherein the bond between C₇ and C₈ is a double bond, which is a compound of formula (I-i), by forming a double bond between C₇ and C₈, in particular via an olefin metathesis reaction, with concomitant cyclization to the macrocycle as outlined in the following reaction scheme:



wherein in the above and following reaction schemes R^8 represents a radical






- 5 (b) converting a compound of formula (I-i) to a compound of formula (I) wherein the link between C7 and C8 in the macrocycle is a single bond, i.e. a compound of formula (I-j):




- 10 by a reduction of the C7-C8 double bond in the compounds of formula (I-j);


- (c) preparing a compound of formula (I) wherein R^1 represents $-NHSO_2R^7$, said compounds being represented by formula (I-k-1), by forming an amide bond between a intermediate (2a) and an sulfonylamine (2b), or preparing a compound of formula (I) wherein R^1 represents $-OR^6$, i.e. a compound (I-k-2), by forming an ester bond between an intermediate (2a) and an alcohol (2c) as outlined in the following scheme wherein G represents a group:
- 15



- 5 (d) preparing a compound of formula (I) wherein R^3 is hydrogen, said compound being represented by (I-l), from a corresponding nitrogen-protected intermediate (3a), wherein PG represents a nitrogen protecting group:



- (e) reacting an intermediate (4a) with an amine (4b) in the presence of a carbamate forming reagent as outlined in the following reaction scheme:



(f) converting compounds of formula (I) into each other by a functional group transformation reaction; or

5

(g) preparing a salt form by reacting the free form of a compound of formula (I) with an acid or a base.

10

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2006/064817

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D401/12 C07D403/12 C07D413/12 A61K31/538 A61P31/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2005/037214 A (INTERMUNE, INC; ARRAY BIOPHARMA INC) 28 April 2005 (2005-04-28) cited in the application Abstract; claims; examples.	1-12
P,A	WO 2005/073216 A (MEDIVIR AB; ROSENQUIST, AASA; THORSTENSSON, FREDRIK; JOHANSSON, PER-OL) 11 August 2005 (2005-08-11) Abstract; claims; examples, e.g. no. 139.	1-12
P,A	WO 2005/073195 A (MEDIVIR AB; ROSENQUIST, AASA; THORSTENSSON, FREDRIK; JOHANSSON, PER-OL) 11 August 2005 (2005-08-11) Abstract; claims; examples, e.g. no. 109.	1-12

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
 "&" document member of the same patent family

Date of the actual completion of the international search

24 October 2006

Date of mailing of the international search report

31/10/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Weisbrod, Thomas

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2006/064817

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 11 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compounds/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2006/064817

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2005037214 A	28-04-2005	AU 2004281780 A1 CA 2540858 A1 EP 1680137 A2	28-04-2005 28-04-2005 19-07-2006
WO 2005073216 A	11-08-2005	AR 048401 A1 AR 047793 A1 AU 2005207815 A1 AU 2005207816 A1 CA 2552317 A1 CA 2552319 A1 WO 2005073195 A2	26-04-2006 22-02-2006 11-08-2005 11-08-2005 11-08-2005 11-08-2005 11-08-2005
WO 2005073195 A	11-08-2005	AR 048401 A1 AR 047793 A1 AU 2005207815 A1 AU 2005207816 A1 CA 2552317 A1 CA 2552319 A1 WO 2005073216 A2	26-04-2006 22-02-2006 11-08-2005 11-08-2005 11-08-2005 11-08-2005 11-08-2005